

**FEATURES:**  
 1. 277  
 2. 273  
 3. 2056  
 4. 2056



SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## **ISOLATED HUMAN TRANSPORTER PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS, AND USES THEREOF**

### **RELATED APPLICATIONS**

The present application is a Continuation-In-Part of U.S. Serial No.09/630,719, filed August 2, 2000 and Continuation-In-Part of U.S. Serial No. 09/765,344.

### **FIELD OF THE INVENTION**

The present invention is in the field of transporter proteins that are related to the sulfate transporter subfamily, recombinant DNA molecules, and protein production. The present invention specifically provides novel peptides and proteins, representing two splice forms of a novel sulfate transporter, that effect ligand transport and nucleic acid molecules encoding such peptide and protein molecules, all of which are useful in the development of human therapeutics and diagnostic compositions and methods.

### **BACKGROUND OF THE INVENTION**

#### Transporters

Transporter proteins regulate many different functions of a cell, including cell proliferation, differentiation, and signaling processes, by regulating the flow of molecules such as ions and macromolecules, into and out of cells. Transporters are found in the plasma membranes of virtually every cell in eukaryotic organisms. Transporters mediate a variety of cellular functions including regulation of membrane potentials and absorption and secretion of molecules and ion across cell membranes. When present in intracellular membranes of the Golgi apparatus and endocytic vesicles, transporters, such as chloride channels, also regulate organelle pH. For a review, see Greger, R. (1988) Annu. Rev. Physiol. 50:111-122.

Transporters are generally classified by structure and the type of mode of action. In addition, transporters are sometimes classified by the molecule type that is transported, for example, sugar transporters, chlorine channels, potassium channels, etc. There may be many classes of channels for transporting a single type of molecule (a detailed review of channel types can be found at Alexander, S.P.H. and J.A. Peters: Receptor and transporter nomenclature

supplement. Trends Pharmacol. Sci., Elsevier, pp. 65-68 (1997) and <http://www-biology.ucsd.edu/~msaier/transport/titlepage2.html>.

### Ion channels

An important type of transporter is the ion channel. Ion channels regulate many different cell proliferation, differentiation, and signaling processes by regulating the flow of ions into and out of cells. Ion channels are found in the plasma membranes of virtually every cell in eukaryotic organisms. Ion channels mediate a variety of cellular functions including regulation of membrane potentials and absorption and secretion of ion across epithelial membranes. When present in intracellular membranes of the Golgi apparatus and endocytic vesicles, ion channels, such as chloride channels, also regulate organelle pH. For a review, see Greger, R. (1988) Annu. Rev. Physiol. 50:111-122.

Ion channels are generally classified by structure and the type of mode of action. For example, extracellular ligand gated channels (ELGs) are comprised of five polypeptide subunits, with each subunit having 4 membrane spanning domains, and are activated by the binding of an extracellular ligand to the channel. In addition, channels are sometimes classified by the ion type that is transported, for example, chlorine channels, potassium channels, etc. There may be many classes of channels for transporting a single type of ion (a detailed review of channel types can be found at Alexander, S.P.H. and J.A. Peters (1997). Receptor and ion channel nomenclature supplement. Trends Pharmacol. Sci., Elsevier, pp. 65-68 and <http://www-biology.ucsd.edu/~msaier/transport/toc.html>.

There are many types of ion channels based on structure. For example, many ion channels fall within one of the following groups: extracellular ligand-gated channels (ELG), intracellular ligand-gated channels (ILG), inward rectifying channels (INR), intercellular (gap junction) channels, and voltage gated channels (VIC). There are additionally recognized other channel families based on ion-type transported, cellular location and drug sensitivity. Detailed information on each of these, their activity, ligand type, ion type, disease association, drugability, and other information pertinent to the present invention, is well known in the art.

### Anion transport proteins

The present invention provides two splice forms of a novel human anion transport protein that shows a particularly high degree of similarity to sulfate transporters. The alternative splice forms are herein referred to as splice forms 1 and 2. Splice form 1 has been previously disclosed by applicant in U.S. application 09/630,719, filed August 2, 2000.



Anion transport proteins in mammalian cells participate in a wide variety of cell and intracellular organelle functions, including regulation of electrical activity, pH, volume, and the transport of osmolites and metabolites. These proteins also have essential physiological roles in the control of immunological responses, cell migration, cell proliferation, and differentiation. Several classes of anion transporters have been characterized with varying molecular structures and mechanisms for mediating anion flux. One of the most prominent anion transporter super-families is the multiple membrane-spanning permeases, which include Na<sup>+</sup>- or H<sup>+</sup>-dependent anion coanion transporters (symporters), anion/anion exchangers (antiporters), and cation-independent anion uniporters. This super-family is also referred to as the Carrier-type anion transporters <<http://www-biology.ucsd.edu/~msaier/transport/titlepage2.html>>. Typically, proteins within this group contain 8-14 hydrophobic alpha-helical peptide segments that allow the protein to reside in the membrane bilayer. These helices also establish the pathway for ion translocation. Both broad-substrate and substrate-specific anion transporters are known: the former type enables multiple anion species (chloride, iodide, sulfate, bromide, etc.) to permeate the bounding membrane, while anion transporters in the latter class restrict ion movement to one chemical species.

Anion transporter genes and gene products are potential causative agents of disease and disease phenotypes may be actuated both by alterations in gene transcription and by mutations in the protein sequence. For example, the down-regulated in adenoma (DRA) gene was originally identified as a gene that was down-regulated in colon tumors. It encodes a protein with anion transporter function that is expressed in the intestinal tract (duodenum, ileum, cecum, distal colon), but not in the esophagus or stomach (Antalis T.M., Reeder J.A., Gotley D.C. *et al.*, *Clin Cancer Res* (1998) Aug;4(8):1857-63; Byeon M.K., Westerman M.A., Maroulakou I.G. *et al.*, *Oncogene* (1996) Jan 18;12(2):387-96). A second illustration of the biomedical significance of anion transporters is found with patients presenting severe hypothyroidism caused by a congenital lack of iodide transport. These individuals do not accumulate-iodide in their thyroids. A single amino acid substitution in the thyroid Na<sup>+</sup>/I<sup>-</sup> symporter, where proline replaced threonine at position 354, has been identified as the cause of this condition in two independent patients (Levy O., Ginter C.S., De la Vieja A. *et al.*, *FEBS Lett* (1998) Jun 5;429(1):36-40). Equally compelling are two well-documented autosomal recessive disorders Pendred syndrome and Diastrophic dysplasia (DTD). Pendred syndrome is the most common form of syndromic deafness and characterized by congenital sensorineural hearing loss and goitre. This disorder has been mapped to chromosome 7 and the gene product causing Pendred syndrome (PDS) has been

identified as a anion transporter for iodide and chloride (Scott D.A., Wang R., Kreman T.M. *et al.*, *Nat Genet* (1999)Apr;21(4):440-3). DTD is a well-characterized osteochondrodysplasia with clinical features including dwarfism, spinal deformation, and specific joint abnormalities. The disease occurs in most populations. The gene has been mapped to distal chromosome 5q and it encodes a sulfate anion transporter (Hastbacka J., de la Chapelle A., Mahtani M.M. *et al.*, *Cell* 1994 Sep 23;78(6):1073-87).

Issued US Patents that demonstrate the utility for this group of protein/DNA molecules include, but are not limited to, 6,054,558 "Compositions and methods for the treatment and diagnosis of cardiovascular disease using rchd534 as a target"; 6,048,709 "Compositions and methods for the treatment and diagnosis of cardiovascular disease"; 6,046,030 "Human LIG-1 homolog (HLIG-1)"; 6,025,160 "Polynucleotide and polypeptide sequences encoding rat mdr1b2 and screening methods thereof"; 6,013,672 "Agonists of metabotropic glutamate receptors and uses thereof"; 6,008,015 "Glycine transporter"; 5,989,825 "Excitatory amino acid transporter gene and uses"; and 5,928,926 "Isolation and cloning of the human ARSA-I gene and uses thereof".

Transporter proteins, particularly members of the sulfate transporter subfamily, are a major target for drug action and development, particularly members that are expressed in the tissue types noted in Figure 1 (e.g. neoplastic cells). Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown transport proteins. The present invention advances the state of the art by providing a previously unidentified human transport protein.

## SUMMARY OF THE INVENTION

The present invention is based in part on the identification of amino acid sequences of human transporter peptides and proteins that are related to the sulfate transporter subfamily, as well as allelic variants and other mammalian orthologs thereof. Specifically, the present invention provides two splice forms of a novel human sulfate transporter protein. These unique peptide sequences, and nucleic acid sequences that encode these peptides, can be used as models for the development of human therapeutic targets, aid in the identification of therapeutic proteins, and serve as targets for the development of human therapeutic agents that modulate transporter activity in cells and tissues that express the transporter. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal

muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas.

### DESCRIPTION OF THE FIGURE SHEETS

FIGURE 1 provides the nucleotide sequences of cDNA molecules that encode splice forms 1 and 2 of the transporter protein of the present invention (splice form 1 = SEQ ID NO:1, splice form 2 = SEQ ID NO:4). In addition, structure and functional information is provided, such as ATG start, stop and tissue distribution, where available, that allows one to readily determine specific uses of the inventions based on these molecular sequences. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas.

FIGURE 2 provides the predicted amino acid sequences of splice forms 1 and 2 of the transporter of the present invention (splice form 1 = SEQ ID NO:2, splice form 2 = SEQ ID NO:5). In addition, structure and functional information such as protein family, function, and modification sites is provided where available, allowing one to readily determine specific uses of the inventions based on these molecular sequences.

FIGURE 3 provides a genomic sequence (SEQ ID NO:3) that spans the gene encoding splice forms 1 and 2 of the transporter protein of the present invention. In addition, structure and functional information, such as intron/exon structure, promoter location, etc., is provided where available, allowing one to readily determine specific uses of the inventions based on this molecular sequence. As illustrated in Figure 3, identified SNP variations include g30344a, a31170g, c16256t, a13376t, t12210c, g12072c, g11922t, -11903a, c10009g, c4519t, a4181g, a20952c, t20987c, g21620a, t21795c, a22753t, g22945a, g23032a, g23738a, t23952g, a24123g, c24527-, c24691t, g25015a, and g25191t. Figure 3 also provides structural information for splice form 2, derived from the Genewise computer program.

## DETAILED DESCRIPTION OF THE INVENTION

### General Description

The present invention is based on the sequencing of the human genome. During the sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a transporter protein or part of a transporter protein and are related to the sulfate transporter subfamily. Specifically, the present invention provides two splice forms of a novel human sulfate transporter. Utilizing these sequences, additional genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of human transporter peptides and proteins that are related to the sulfate transporter subfamily, nucleic acid sequences in the form of transcript sequences, cDNA sequences and/or genomic sequences that encode these transporter peptides and proteins, nucleic acid variation (allelic information), tissue distribution of expression, and information about the closest art known protein/peptide/domain that has structural or sequence homology to the transporter of the present invention.

In addition to being previously unknown, the peptides that are provided in the present invention are selected based on their ability to be used for the development of commercially important products and services. Specifically, the present peptides are selected based on homology and/or structural relatedness to known transporter proteins of the sulfate transporter subfamily and the expression pattern observed. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas.. The art has clearly established the commercial importance of members of this family of proteins and proteins that have expression patterns similar to that of the present gene. Some of the more specific features of the peptides of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for each of the known sulfate transporter family or subfamily of transporter proteins.

## Specific Embodiments

### Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being members of the transporter family of proteins and are related to the sulfate transporter subfamily (protein sequences are provided in Figure 2, cDNA sequences are provided in Figures 1 and genomic sequences are provided in Figure 3). Specifically, the present invention provides two splice forms of a novel human sulfate transporter. The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the transporter peptides of the present invention, transporter peptides, or peptides/proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprising the amino acid sequences of the transporter peptides disclosed in the Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The peptides of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components (the features of an isolated nucleic acid molecule is discussed below).

In some uses, "substantially free of cellular material" includes preparations of the peptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical

precursors or other chemicals" includes preparations of the transporter peptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

The isolated transporter peptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. For example, a nucleic acid molecule encoding the transporter peptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

Accordingly, the present invention provides proteins that consist of the amino acid sequences provided in Figure 2 (SEQ ID NOS:2 and 5), for example, proteins encoded by the cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NOS:1 and 4) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid sequence when the amino acid sequence is the final amino acid sequence of the protein.

The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NOS:2 and 5), for example, proteins encoded by the cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NOS:1 and 4) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about 1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NOS:2 and 5), for example, proteins encoded by the cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NOS:1 and 4) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein comprises an amino acid sequence when the amino acid

sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the transporter peptides of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

The transporter peptides of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a transporter peptide operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the transporter peptide. "Operatively linked" indicates that the transporter peptide and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the transporter peptide.

In some uses, the fusion protein does not affect the activity of the transporter peptide *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant transporter peptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A transporter peptide-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the transporter peptide.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the proteins of the present invention, such as naturally occurring mature

forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogs of the peptides. Such variants can readily be generated using art-known techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the transporter peptides of the present invention. The degree of homology/identity present will be based primarily on whether the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.

To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of a reference sequence is aligned for comparison purposes. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part 1*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm



which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.* 12(1):387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Myers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

Full-length pre-processed forms, as well as mature processed forms, of proteins that comprise one of the peptides of the present invention can readily be identified as having complete sequence identity to one of the transporter peptides of the present invention as well as being encoded by the same genetic locus as the transporter peptide provided herein. RH panel mapping shows the gene encoding the transporter proteins of the present invention is found on chromosome 17 near markers SHGC-56719 and SHGC-58932 (LOD=15.9 and 15.68).

Allelic variants of a transporter peptide can readily be identified as being a human protein having a high degree (significant) of sequence homology/identity to at least a portion of the transporter peptide as well as being encoded by the same genetic locus as the transporter peptide provided herein. Genetic locus can readily be determined based on the genomic information

provided in Figure 3, such as the genomic sequence mapped to the reference human. RH panel mapping shows the gene encoding the transporter proteins of the present invention is found on chromosome 17 near markers SHGC-56719 and SHGC-58932 (LOD=15.9 and 15.68). As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a transporter peptide encoding nucleic acid molecule under stringent conditions as more fully described below.

Figure 3 provides information on SNPs that have been found in the gene encoding the transporter proteins of the present invention. The following variations were seen: g30344a, a31170g, c16256t, a13376t, t12210c, g12072c, g11922t, -11903a, c10009g, c4519t, a4181g, a20952c, t20987c, g21620a, t21795c, a22753t, g22945a, g23032a, g23738a, t23952g, a24123g, c24527-, c24691t, g25015a, and g25191t.

Paralogs of a transporter peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the transporter peptide, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are typically at least about 60% or greater, and more typically at least about 70% or greater homology through a given region or domain. Such paralogs will be encoded by a nucleic acid sequence that will hybridize to a transporter peptide encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a transporter peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the transporter peptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a transporter peptide encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the transporter peptides of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the transporter peptide. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those that substitute a given amino acid in a transporter peptide by another amino acid of like

characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant transporter peptides can be fully functional or can lack function in one or more activities, e.g. ability to bind ligand, ability to transport ligand, ability to mediate signaling, etc. Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Figure 2 provides the result of protein analysis and can be used to identify critical domains/regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 244:1081-1085 (1989)), particularly using the results provided in Figure 2. The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as transporter activity or in assays such as an *in vitro* proliferative activity. Sites that are critical for binding partner/substrate binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.* *Science* 255:306-312 (1992)).

The present invention further provides fragments of the transporter peptides, in addition to proteins and peptides that comprise and consist of such fragments, particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that may be disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16, or more contiguous amino acid residues from a transporter peptide. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the transporter peptide or could be chosen for the

ability to perform a function, e.g. bind a substrate or act as an immunogen. Particularly important fragments are biologically active fragments, peptides that are, for example, about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the transporter peptide, e.g., active site, a transmembrane domain or a substrate-binding domain. Further, possible fragments include, but are not limited to, domain or motif containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well known and readily available to those of skill in the art (e.g., PROSITE analysis). The results of one such analysis are provided in Figure 2.

Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in transporter peptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the transporter peptides of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature transporter peptide is fused with another compound, such as a compound to increase the half-life of the transporter peptide (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature transporter peptide, such as a leader or secretory sequence or a sequence for purification of the mature transporter peptide or a pro-protein sequence.

#### Protein/Peptide Uses

The proteins of the present invention can be used in substantial and specific assays related to the functional information provided in the Figures; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its binding partner or ligand) in biological fluids; and as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein or ligand (such as, for example, in a transporter-effector protein interaction or transporter-ligand interaction), the protein can be used to identify the binding partner/ligand so as to develop a system to identify inhibitors of the binding interaction. Any or all of these uses are capable of being developed into reagent grade or kit format for commercialization as commercial products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

The potential uses of the peptides of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, transporters isolated from humans and their human/mammalian orthologs serve as targets for identifying agents for use in mammalian therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the transporter. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas.

Additionally, BLAST hits to ESTs derived from tissue specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen. A large percentage of pharmaceutical agents are being developed that modulate the activity of transporter proteins, particularly members of the sulfate transporter subfamily (see Background of the Invention). The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the expression information provided in Figure 1. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemia B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. Such uses can readily be determined using the information provided herein, that known in the art and routine experimentation.

The proteins of the present invention (including variants and fragments that may have been disclosed prior to the present invention) are useful for biological assays related to transporters that are related to members of the sulfate transporter subfamily. Such assays involve any of the known transporter functions or activities or properties useful for diagnosis and treatment of transporter-related conditions that are specific for the subfamily of transporters that the one of the present invention belongs to, particularly in cells and tissues that express the transporter. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas. Additionally, BLAST hits to ESTs derived from tissue specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen. The proteins of the present invention are also useful in drug screening assays, in cell-based or cell-free systems ((Hodgson, Bio/technology, 1992, Sept 10(9);973-80). Cell-based systems can be native, i.e., cells that normally express the transporter, as a biopsy or expanded in cell culture. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated

tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the transporter protein.

The polypeptides can be used to identify compounds that modulate transporter activity of the protein in its natural state or an altered form that causes a specific disease or pathology associated with the transporter. Both the transporters of the present invention and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to the transporter. These compounds can be further screened against a functional transporter to determine the effect of the compound on the transporter activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the transporter to a desired degree.

Further, the proteins of the present invention can be used to screen a compound for the ability to stimulate or inhibit interaction between the transporter protein and a molecule that normally interacts with the transporter protein, e.g. a substrate or a component of the signal pathway that the transporter protein normally interacts (for example, another transporter). Such assays typically include the steps of combining the transporter protein with a candidate compound under conditions that allow the transporter protein, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the transporter protein and the target, such as any of the associated effects of signal transduction such as changes in membrane potential, protein phosphorylation, cAMP turnover, and adenylate cyclase activation, etc.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')<sub>2</sub>, Fab expression library

fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries).

One candidate compound is a soluble fragment of the receptor that competes for ligand binding. Other candidate compounds include mutant transporters or appropriate fragments containing mutations that affect transporter function and thus compete for ligand. Accordingly, a fragment that competes for ligand, for example with a higher affinity, or a fragment that binds ligand but does not allow release, is encompassed by the invention.

The invention further includes other end point assays to identify compounds that modulate (stimulate or inhibit) transporter activity. The assays typically involve an assay of events in the signal transduction pathway that indicate transporter activity. Thus, the transport of a ligand, change in cell membrane potential, activation of a protein, a change in the expression of genes that are up- or down-regulated in response to the transporter protein dependent signal cascade can be assayed.

Any of the biological or biochemical functions mediated by the transporter can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art or that can be readily identified using the information provided in the Figures, particularly Figure 2. Specifically, a biological function of a cell or tissues that expresses the transporter can be assayed. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas. Additionally, BLAST hits to ESTs derived from tissue specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen.

Binding and/or activating compounds can also be screened by using chimeric transporter proteins in which the amino terminal extracellular domain, or parts thereof, the entire transmembrane domain or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by heterologous domains or subregions. For example, a ligand-binding region can be used that interacts with a different ligand than that which is recognized by the native transporter. Accordingly, a different set of signal transduction components is available as an end-



point assay for activation. This allows for assays to be performed in other than the specific host cell from which the transporter is derived.

The proteins of the present invention are also useful in competition binding assays in methods designed to discover compounds that interact with the transporter (e.g. binding partners and/or ligands). Thus, a compound is exposed to a transporter polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide. Soluble transporter polypeptide is also added to the mixture. If the test compound interacts with the soluble transporter polypeptide, it decreases the amount of complex formed or activity from the transporter target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the transporter. Thus, the soluble polypeptide that competes with the target transporter region is designed to contain peptide sequences corresponding to the region of interest.

To perform cell free drug screening assays, it is sometimes desirable to immobilize either the transporter protein, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g.,  $^{35}\text{S}$ -labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of transporter-binding protein found in the bead fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a transporter-binding protein and a candidate compound are incubated in the transporter protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the

GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the transporter protein target molecule, or which are reactive with transporter protein and compete with the target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the transporters of the present invention can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal or other model system. Such model systems are well known in the art and can readily be employed in this context.

Modulators of transporter protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the transporter pathway, by treating cells or tissues that express the transporter. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. These methods of treatment include the steps of administering a modulator of transporter activity in a pharmaceutical composition to a subject in need of such treatment, the modulator being identified as described herein.

In yet another aspect of the invention, the transporter proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with the transporter and are involved in transporter activity. Such transporter-binding proteins are also likely to be involved in the propagation of signals by the transporter proteins or transporter targets as, for example, downstream elements of a transporter-mediated signaling pathway. Alternatively, such transporter-binding proteins are likely to be transporter inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a transporter protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an

unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a transporter-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the transporter protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a transporter-modulating agent, an antisense transporter nucleic acid molecule, a transporter-specific antibody, or a transporter-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

The transporter proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. The method involves contacting a biological sample with a compound capable of interacting with the transporter protein such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The peptides of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered transporter activity in cell-based or cell-free assay, alteration in ligand or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

*In vitro* techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody or other types of detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes effects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from

standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the transporter protein in which one or more of the transporter functions in one population is different from those in another population. The peptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other ligand-binding regions that are more or less active in ligand binding, and transporter activation. Accordingly, ligand dosage would necessarily be modified to maximize the therapeutic effect within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. Accordingly, methods for treatment include the use of the transporter protein or fragments.

### Antibodies

The invention also provides antibodies that selectively bind to one of the peptides of the present invention, a protein comprising such a peptide, as well as variants and fragments thereof. As used herein, an antibody selectively binds a target peptide when it binds the target peptide and does not significantly bind to unrelated proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still selective despite some degree of cross-reactivity.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')<sub>2</sub>, and Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, *Antibodies*, Cold Spring Harbor Press, (1989).

In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the transporter proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or transporter/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence alignment can be used to identify conserved and unique sequence fragments.

An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such fragments can be selected on a physical property, such as fragments correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2).

Detection on an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$  or  $^3\text{H}$ .

### Antibody Uses

The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues in an organism and over the course of normal development. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas. Additionally, BLAST hits to ESTs derived from tissue specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Also, such antibodies can be used to assess abnormal tissue distribution or abnormal expression during development or progression of a biological condition. Antibody detection of circulating fragments of the full length protein can be used to identify turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function. When a disorder is caused by an inappropriate tissue distribution, developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. If a disorder is characterized by a specific mutation in the protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Independent lines of evidence show expression of splice

form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the protein or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. Thus, where a specific protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the transporter peptide to a binding partner such as a ligand or protein binding partner. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the peptides activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount



of protein in the sample; means for comparing the amount of protein in the sample with a standard; and instructions for use. Such a kit can be supplied to detect a single protein or epitope or can be configured to detect one of a multitude of epitopes, such as in an antibody detection array. Arrays are described in detail below for nucleic acid arrays and similar methods have been developed for antibody arrays.

### Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a transporter peptide or protein of the present invention (cDNA, transcript and genomic sequence). Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide sequence that encodes one of the transporter peptides of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.

As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences that naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide sequences, for example up to about 5KB, 4KB, 3KB, 2KB, or 1KB or less, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations described herein such as recombinant expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a transcript/cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NOS:1 and 4, cDNA/transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NOS:2 and 5. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide sequence is the complete nucleotide sequence of the nucleic acid molecule.

The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NOS:1 and 4, cDNA/transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NOS:2 and 5. A nucleic acid molecule consists essentially of a nucleotide sequence when such a nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NOS:1 and 4, cDNA/transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NOS:2 and 5. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional nucleotides or can comprise several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, humans genomic sequence (Figure 3) and cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when the mature form

has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the transporter peptide alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides nucleic acid molecules that encode fragments of the peptides of the present invention as well as nucleic acid molecules that encode obvious variants of the transporter proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

The present invention further provides non-coding fragments of the nucleic acid molecules provided in Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences and gene termination sequences. Such fragments are useful in controlling heterologous gene expression and in

developing screens to identify gene-modulating agents. A promoter can readily be identified as being 5' to the ATG start site in the genomic sequence provided in Figure 3.

A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could be at least 30, 40, 50, 100, 250 or 500 nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope bearing regions of the peptide, or can be useful as DNA probes and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further, primers can be used in PCR reactions to clone specific regions of gene.

A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 20, 25, 40, 50 or more consecutive nucleotides.

Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. Allelic variants can readily be determined by genetic locus of the encoding gene. RH panel mapping shows the gene encoding the transporter proteins of the present invention is found on chromosome 17 near markers SHGC-56719 and SHGC-58932 (LOD=15.9 and 15.68).

Figure 3 provides information on SNPs that have been found in the gene encoding the transporter proteins of the present invention. The following variations were seen: g30344a, a31170g, c16256t, a13376t, t12210c, g12072c, g11922t, -11903a, c10009g, c4519t, a4181g, a20952c, t20987c, g21620a, t21795c, a22753t, g22945a, g23032a, g23738a, t23952g, a24123g, c24527-, c24691t, g25015a, and g25191t.

As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The conditions can be such that sequences at least about 60%, at least about 70%, or at least about 80% or more

homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. One example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65C. Examples of moderate to low stringency hybridization conditions are well known in the art.

### Nucleic Acid Molecule Uses

The nucleic acid molecules of the present invention are useful for probes, primers, chemical intermediates, and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the same or related peptides shown in Figure 2. As illustrated in Figure 3, identified SNP variations include g30344a, a31170g, c16256t, a13376t, t12210c, g12072c, g11922t, -11903a, c10009g, c4519t, a4181g, a20952c, t20987c, g21620a, t21795c, a22753t, g22945a, g23032a, g23738a, t23952g, a24123g, c24527-, c24691t, g25015a, and g25191t.

The probe can correspond to any sequence along the entire length of the nucleic acid molecules provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be construed as encompassing fragments disclosed prior to the present invention.

The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.

The nucleic acid molecules are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

The nucleic acid molecules are also useful as probes for determining the chromosomal positions of the nucleic acid molecules by means of *in situ* hybridization methods. RH panel

mapping shows the gene encoding the transporter proteins of the present invention is found on chromosome 17 near markers SHGC-56719 and SHGC-58932 (LOD=15.9 and 15.68).

The nucleic acid molecules are also useful in making vectors containing the gene regulatory regions of the nucleic acid molecules of the present invention.

The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful as hybridization probes for determining the presence, level, form and distribution of nucleic acid expression. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas. Additionally, BLAST hits to ESTs derived from tissue specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen.

Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic acid whose level is determined can be DNA or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or decrease in transporter protein expression relative to normal results.

*In vitro* techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA include Southern hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a transporter protein, such as by measuring a level of a transporter-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if a transporter gene

has been mutated. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas. Additionally, BLAST hits to ESTs derived from tissue specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen.

Nucleic acid expression assays are useful for drug screening to identify compounds that modulate transporter nucleic acid expression.

The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the transporter gene, particularly biological and pathological processes that are mediated by the transporter in cells and tissues that express it. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemia B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas. The method typically includes assaying the ability of the compound to modulate the expression of the transporter nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired transporter nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the transporter nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

The assay for transporter nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds involved in the signal pathway. Further, the expression of genes that are up- or down-regulated in response to the transporter protein signal pathway can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of transporter gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of transporter mRNA in the presence of the candidate compound is compared to the level of expression of transporter mRNA in the absence of the candidate compound. The candidate

compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate transporter nucleic acid expression in cells and tissues that express the transporter. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas. Additionally, BLAST hits to ESTs derived from tissue specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) of nucleic acid expression.

Alternatively, a modulator for transporter nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits the transporter nucleic acid expression in the cells and tissues that express the protein. Independent lines of evidence show expression of splice form 1 in fetal tissues such as brain and kidney; differentiated tissues such as brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas; and neoplastic tissues such as germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendrogliomas, chronic lymphocytic leukemic B-cells, glioblastomas, and well-differentiated endometrial adenocarcinomas.

The nucleic acid molecules are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of the transporter gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a



physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in transporter nucleic acid expression, and particularly in qualitative changes that lead to pathology. The nucleic acid molecules can be used to detect mutations in transporter genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally occurring genetic mutations in the transporter gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of the transporter gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a transporter protein.

Individuals carrying mutations in the transporter gene can be detected at the nucleic acid level by a variety of techniques. Figure 3 provides information on SNPs that have been found in the gene encoding the transporter proteins of the present invention. The following variations were seen: g30344a, a31170g, c16256t, a13376t, t12210c, g12072c, g11922t, -11903a, c10009g, c4519t, a4181g, a20952c, t20987c, g21620a, t21795c, a22753t, g22945a, g23032a, g23738a, t23952g, a24123g, c24527-, c24691t, g25015a, and g25191t. RH panel mapping shows the gene encoding the transporter proteins of the present invention is found on chromosome 17 near markers SHGC-56719 and SHGC-58932 (LOD=15.9 and 15.68). Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364 (1994)), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.*, *Nucleic Acids Res.* 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample,

contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

Alternatively, mutations in a transporter gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant transporter gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve, C.W., (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.*, *Adv. Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144 (1993); and Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include selective oligonucleotide hybridization, selective amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the

individual's response to a compound used for treatment (pharmacogenomic relationship). Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the transporter gene in an individual in order to select an appropriate compound or dosage regimen for treatment. Figure 3 provides information on SNPs that have been found in the gene encoding the transporter proteins of the present invention. The following variations were seen: g30344a, a31170g, c16256t, a13376t, t12210c, g12072c, g11922t, -11903a, c10009g, c4519t, a4181g, a20952c, t20987c, g21620a, t21795c, a22753t, g22945a, g23032a, g23738a, t23952g, a24123g, c24527-, c24691t, g25015a, and g25191t.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control transporter gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of transporter protein. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into transporter protein.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of transporter nucleic acid. Accordingly, these molecules can treat a disorder characterized by abnormal or undesired transporter nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the transporter protein, such as ligand binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in transporter gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired transporter protein to treat the individual.

The invention also encompasses kits for detecting the presence of a transporter nucleic acid in a biological sample. Screening of tissue specific cDNA libraries for cDNA retrieval showed expression of splice form 1 in fetal brain, brain, pituitary gland, heart, leukocytes, kidney, liver, thyroid, lung, placenta, skeletal muscle, fetal kidney, small intestine, prostate, testis, adrenal gland, bone marrow, and pancreas. Additionally, BLAST hits to ESTs derived from tissue

specific libraries shows expression of splice form 1 in germ cell tumors, lung carcinoid tissue, brain anaplastic oligodendroglioma, chronic lymphocytic leukemia B-cells, infant brain (73 days post natal), placenta, glioblastoma, well-differentiated endometrial adenocarcinoma, and fetal liver and spleen. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting transporter nucleic acid in a biological sample; means for determining the amount of transporter nucleic acid in the sample; and means for comparing the amount of transporter nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect transporter protein mRNA or DNA.

### Nucleic Acid Arrays

The present invention further provides nucleic acid detection kits, such as arrays or microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1, 3, and 4).

As used herein "Arrays" or "Microarrays" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared and used according to the methods described in US Patent 5,837,832, Chee *et al.*, PCT application W095/11995 (Chee *et al.*), Lockhart, D. J. *et al.* (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. *et al.* (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown *et al.*, US Patent No. 5,807,522.

The microarray or detection kit is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray or detection kit, it may be preferable to use oligonucleotides that are only 7-20 nucleotides in length. The microarray or detection kit may contain oligonucleotides that cover the known 5', or 3', sequence, sequential oligonucleotides that cover the full length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray or detection kit may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray or detection kit, the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm which starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray or detection kit. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler *et al.*) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray or detection kit, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray or detection kit so that the probe sequences hybridize to complementary oligonucleotides of the microarray or detection kit. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative

abundance of each oligonucleotide sequence on the microarray or detection kit. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large-scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression of the transporter proteins/peptides of the present invention. In detail, such methods comprise incubating a test sample with one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention and or alleles of the transporter gene of the present invention. Figure 3 provides information on SNPs that have been found in the gene encoding the transporter proteins of the present invention. The following variations were seen: g30344a, a31170g, c16256t, a13376t, t12210c, g12072c, g11922t, -11903a, c10009g, c4519t, a4181g, a20952c, t20987c, g21620a, t21795c, a22753t, g22945a, g23032a, g23738a, t23952g, a24123g, c24527-, c24691t, g25015a, and g25191t.

Conditions for incubating a nucleic acid molecule with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the Human genome disclosed herein. Examples of such assays can be found in Chard, T, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the Human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid.

In detail, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified transporter gene of the present invention can be routinely identified using the sequence information disclosed herein can be readily incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

#### Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in procaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the nucleic acid molecules from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage  $\lambda$ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses,



papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and eukaryotic hosts are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterotrinsporter. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann

*et al.*, *Gene* 69:301-315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)).

Recombinant protein expression can be maximized in host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the nucleic acid molecule sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters

described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

Where secretion of the peptide is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as transporters, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

Where the peptide is not secreted into the medium, which is typically the case with transporters, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the peptides described herein, the peptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may include an initial modified methionine in some cases as a result of a host-mediated process.

#### Uses of vectors and host cells

The recombinant host cells expressing the peptides described herein have a variety of uses. First, the cells are useful for producing a transporter protein or peptide that can be further purified to produce desired amounts of transporter protein or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the transporter protein or transporter protein fragments, such as those described above as well as other formats known in the art. Thus, a recombinant host cell expressing a native transporter protein is useful for assaying compounds that stimulate or inhibit transporter protein function.

Host cells are also useful for identifying transporter protein mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant transporter protein (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native transporter protein.

Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse,

in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA that is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a transporter protein and identifying and evaluating modulators of transporter protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the transporter protein nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the transporter protein to particular cells.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al.* *PNAS* 89:6232-6236 (1992). Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al.* *Science* 251:1351-1355 (1991)). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein

is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G<sub>0</sub> phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein are useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect ligand binding, transporter protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* transporter protein function, including ligand interaction, the effect of specific mutant transporter proteins on transporter protein function and ligand interaction, and the effect of chimeric transporter proteins. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more transporter protein functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

### Claims

That which is claimed is:

1. An isolated peptide consisting of an amino acid sequence selected from the group consisting of:

- (a) an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5;
- (b) an amino acid sequence of an allelic variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4;
- (c) an amino acid sequence of an ortholog of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4; and
- (d) a fragment of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said fragment comprises at least 10 contiguous amino acids.

2. An isolated peptide comprising an amino acid sequence selected from the group consisting of:

- (a) an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5;
- (b) an amino acid sequence of an allelic variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4;
- (c) an amino acid sequence of an ortholog of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4; and
- (d) a fragment of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said fragment comprises at least 10 contiguous amino acids.

3. An isolated antibody that selectively binds to a peptide of claim 2.
4. An isolated nucleic acid molecule consisting of a nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5;
  - (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4;
  - (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4;
  - (d) a nucleotide sequence that encodes a fragment of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said fragment comprises at least 10 contiguous amino acids; and
  - (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).
5. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5;
  - (b) a nucleotide sequence that encodes an allelic variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4;
  - (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4;



(d) a nucleotide sequence that encodes a fragment of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5, wherein said fragment comprises at least 10 contiguous amino acids; and

(e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

6. A gene chip comprising a nucleic acid molecule of claim 5.
7. A transgenic non-human animal comprising a nucleic acid molecule of claim 5.
8. A nucleic acid vector comprising a nucleic acid molecule of claim 5.
9. A host cell containing the vector of claim 8.
10. A method for producing any of the peptides of claim 1 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
11. A method for producing any of the peptides of claim 2 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
12. A method for detecting the presence of any of the peptides of claim 2 in a sample, said method comprising contacting said sample with a detection agent that specifically allows detection of the presence of the peptide in the sample and then detecting the presence of the peptide.
13. A method for detecting the presence of a nucleic acid molecule of claim 5 in a sample, said method comprising contacting the sample with an oligonucleotide that hybridizes to said nucleic acid molecule under stringent conditions and determining whether the oligonucleotide binds to said nucleic acid molecule in the sample.

14. A method for identifying a modulator of a peptide of claim 2, said method comprising contacting said peptide with an agent and determining if said agent has modulated the function or activity of said peptide.
15. The method of claim 14, wherein said agent is administered to a host cell comprising an expression vector that expresses said peptide.
16. A method for identifying an agent that binds to any of the peptides of claim 2, said method comprising contacting the peptide with an agent and assaying the contacted mixture to determine whether a complex is formed with the agent bound to the peptide.
17. A pharmaceutical composition comprising an agent identified by the method of claim 16 and a pharmaceutically acceptable carrier therefor.
18. A method for treating a disease or condition mediated by a human transporter protein, said method comprising administering to a patient a pharmaceutically effective amount of an agent identified by the method of claim 16.
19. A method for identifying a modulator of the expression of a peptide of claim 2, said method comprising contacting a cell expressing said peptide with an agent, and determining if said agent has modulated the expression of said peptide.

20. An isolated human transporter peptide having an amino acid sequence that shares at least 70% homology with an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5.

21. A peptide according to claim 20 that shares at least 90 percent homology with an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 5.

22. An isolated nucleic acid molecule encoding a human transporter peptide, said nucleic acid molecule sharing at least 80 percent homology with a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4.

23. A nucleic acid molecule according to claim 22 that shares at least 90 percent homology with a nucleic acid molecule selected from the group consisting of SEQ ID NOS:1, 3, and 4.

## SPLICE FORM 1:

```

1 AACAGCACGA GGGCGGACCC AGCTGTGGCG ACGCCAGGAG ACCCCAAGCT
51 GCATCGCCGA GTGGAAGCAA CTAGAACTCC AGGGCTGTGA AAGCCACAGG
101 TGGGGGGCTGA GCGAGGCGTG GCCTCAGGAG CGGAGGACCC CCCACTCTC
151 CCTCGAGCGC CGCAGTCCAC CGTAGCGGGT GGAGCCCGCC TTGGTGCGCA
201 GTTGGAAAAC CTCGGAGCCC CGCTGGATCT CCTGGCTGCC ACCCGCACCC
251 CCCGCCAGCC TACGCCCCAC CGTAGAGATG CCTTCTTCGG TGACGGCGCT
301 GGGTCAGGCC AGGTCTCTG GCGCCGGGAT GCGCCCGAGC GCCTGCTGCT
351 GCTCCCTGCG GGCCTGCGAG AGGAGGCTGC CCATCTGGC GTGGCTGCCC
401 AGCTACTCCC TGCAGTGGCT GAAGATGGAT TTCGTGCGCC GCCTCTCAGT
451 TGGCCTCACT GCCATTCCCC AGGCGCTGGC CTATGCTGAA GTGGCTGGAC
501 TCCCGCCCCA GTATGGCCTC TACTCTGCCT TCATGGGCTG CTTGCTGTAT
551 TTCTTCTTGG GCACCTCCCG GGATGTGACT CTGGGCCCCA CCGCATTAT
601 GTCCCTCCTG GTCTCCTTCT ACACCTTCCA TGAGCCCGCC TACGCTGTGC
651 TGCTGGCCTT CCTGTCCGGC TGCATCCAGC TGGCCATGGG GGTCTGCGT
701 TTGGGGTTCC TGCTGGACTT CATTTCTTAC CCCGTCATTA AAGGCTTCAC
751 CTCTGCTGCT GCCGTCACCA TCGGCTTTGG ACAGATCAAG AACCTGCTGG
801 GACTACAGAA CATCCCCAGG CCGTCTTCC TGCAGGTGTA CCACACCTTC
851 CTCAGGATTG CAGAGACCAG GGTAGGTGAC GCGCTCCTGG GGCTGGTCTG
901 CATGTGCTG CTGCTGGTGC TGAAGCTGAT GCGGGACCAC GTGCCTCCCG
951 TCCACCCCGA GATGCCCCCT GGTGTGCGGC TCAGCCGTGG GCTGGTCTGG
1001 GCTGCCACGA CAGCTCGCAA CGCCCTGGTG GTCTCCTTCG CAGCCCTGGT
1051 TGCGTACTCC TTCGAGGTGA CTGGATACCA GCCTTTCATC CTAACAGGGG
1101 AGACAGCTGA GGGGCTCCCT CCAGTCCGGA TCCCGCCCTT CTCAGTGACC
1151 ACAGCCAACG GGACGATCTC CTTACCCGAG ATGGTGCAGG ACATGGGAGC
1201 CGGGCTGGCC GTGGTGCCCC TGATGGGCTT CCTGGAGAGC ATTGCGGTGG
1251 CCAAAGCCTT CGCATCTCAG AATAATTACC GCATCGATGC CAACCAGGAG
1301 CTGCTGGCCA TCGGTCTCAC CAACATGTTG GGCTCCCTCG TCTCCTCCTA
1351 CCCGTCACA GGCAGCTTTG GACGGACAGC CGTGAACGCT CAGTCGGGGG
1401 TGTGACCCCC GCGGGGGGGC CTGGTGACGG GAGTGCTGGT GCTGCTGTCT
1451 CTGGACTACC TGACCTCACT GTTCTACTAC ATCCCCAAGT CTGCCCTGGC
1501 TGCCGTCATC ATCATGGCCG TGGCCCGCT GTTCGACACC AAGATCTTCA
1551 GGACGCTCTG GCGTGTTAAG AGGCTGGACC TGCTGCCCT GTGCGTGACC
1601 TTCCTGCTGT GCTTCTGGGA GGTGCAGTAC GGCATCCTGG CCGGGGCCCT
1651 GGTGTCTCTG CTCATGCTCC TGCACTCTGC AGCCAGGCCT GAGACCAAGG
1701 TGTGAGAGG GCGGTTCTG GTCTGCAGC CGGCCAGCGG CCTGTCTTC
1751 CCTGCCATGG AGGCTCTGCG GGAGGAGATC CTAAGCCGG CCCTGGAAGT
1801 GTCCCGGCCA CGCTGCCTGG TCCTGGAGTG CACCCATGTC TGCAGCATCG
1851 ACTACACTGT GGTGCTGGGA CTCGGCGAGC TCCTCCAGGA CTCCAGAAG
1901 CAGGCGCTCG CCCTGGCCTT TGTGGGCTG CAGGTCCCG TTCTCCGTGT
1951 CCTGCTGTCC GCTGACCTGA AGGGTTCCA GTACTTCTCT ACCCTGGAAG
2001 AAGCAGAGAA GCACCTGAGG CAGGAGCCAG GGACCCAGCC CTACAACATC
2051 AGAGAAGACT CCATTCTGGA CCAAAGGTT GCCCTGCTCA AGGCATAATG
2101 GGGCCACCCG TGGGCATCCA CAGTTTGCA GGTGTTCCGG AAGGTTCTTG
2151 TCACTGTGAT TGGATGCTGG ATGCCGCTG ATAGACATG TGGCCTGGCT
2201 GAGAAACCCC TGAGCAGGTA ACCCAGGGAA GAGAAGGAAG CCAGGCCTGG
2251 AGGTCCACGG CAGTGGGAGT GGGGCTCACT GGCTTCCTGT GGGATGACTG
2301 GAAATGACC TCGCTGCTGT TCCCTGGCAT GACCTCTTT GGAAGAGTGG
2351 TTTGGAGAGA GCCTTCTAGA ATGACAGACT GTGCGAGGAA GCAGGGGCAG
2401 GGGTTTCCAG CCCGGGCTGT GCGAGGCATC CTGGGGCTGG CAGCACCTTC
2451 CCGGCTCACC AGTGCCACCT GCGGGGGAGG GACGGGGCAG GCAGGAGTCT
2501 GGGAGGCGGG TCCGCTCCTC TTGTCTGCGG CATCTGTGCT CTCCGAGAGA
2551 AAACCAAGGT GTGTCAAATG ACGTCAAGTC TCTATTTAAA AATAATTTTG
2601 TGTTTTCTAA ATGGAAAAAG TGATAGCTTT GGTGATTTTG TAAAAGTCAT
2651 AAATGCTTAT TGTAAAAAAT ACAGGAAACC ACCCTCACC CTGTCCACTT
2701 GGGTGATCAT TCCAGACCCC TCCCCAAACA TGCATATGTA CCTGTCCGTC
2751 AGTGTGTGGA TGTATGTTA CAGTTCTACA TAAATGGGAT CATTTTATAC
2801 ATGGTGCTCT GGAACCCACA TTTTTCATGC AGTCATTTGC AGTGAATTAT
2851 TTATTGTGAT AATAAATAGC ATTAGAATAC AAGAAAAAAA AAAAAAAA
2901 AAAAAAAA AAAAAAAA (SEQ ID NO:1)

```

## FEATURES:

5' UTR: 1-277

Start: 278

Stop: 2096

3' UTR: 2096-2919

## SPLICE FORM 2:

```

1   GCTGTGGCGA  CGCCAGGAGA  CCCCAGCTG  CATCGCCGAG  TGAAGCAAC
51  TAGAACTCCA  GGGCTGTGAA  AGCCACAGGT  GGGGGCTGAG  CGAGGCGTGG
101 CCTCAGGAGC  GGAGGACCCC  CCCACTCTCC  CTCGAGCGCC  GCAGTCCACC
151 GTAGCGGGTG  GAGCCCGCCT  TGGTGCAGAG  TTGGAAAACC  TCGGAGCCCC
201 GCTGGATCTC  CTGGCTGCCA  CCCGCACCCC  CCGCCAGCCT  ACGCCCCACC
251 GTAGAGATGC  CTTCTTCGGT  GACGGCGCTG  GGTGAGGCCA  GGTCTCTGG
301 CCCCAGGATG  GCCCCGAGCG  CCTGTGCTG  CTCCCCTGCG  GCCCTGCAGA
351 GGAGGCTGCC  CATCTGGCG  TGGCTGCCCC  GCTACTCCCT  GCAGTGGCTG
401 AAGATGGATT  TCGTCGCCG  CCTCTCAGTT  GGCCTCACTG  CCATTCCCCA
451 GGCCTGGCC  TATGCTGAAG  TGGCTGGACT  CCGCCCCAG  TATGGCCTCT
501 ACTCTGCCTT  CATGGGCTGC  TTCGTGTATT  TCTTCTGGG  CACCTCCCGG
551 GATGTGACTC  TGGGCCCCAC  CGCCATTATG  TCCCTCCTGG  TCTCCTTCTA
601 CACCTTCCAT  GAGCCCGCCT  ACGCTGTGCT  GCTGGCCTTC  CTGTCCGGCT
651 GCATCCAGCT  GGCCATGGGG  GTCCTGCGTT  TGGCTCACAT  CTCCCCTCAT
701 CCTCTGGGAC  TGGGTGGAGC  CGGACCAGC  TCGATGTCCC  CTCTTGGCTG
751 GCCAGGGTTC  CTGCTGGACT  TCATTTCCTA  CCCCCTCATT  AAAGGCTTCA
801 CCTCTGCTGC  TGCCGTCACC  ATCGGCTTTG  GACAGATCAA  GAACCTGCTG
851 GGAATACAGA  ACATCCCCAG  GCCGTCTTTC  CTGCAGGTGT  ACCACACCTT
901 CCTCAGGATT  GCAGAGACCA  GGGTAGGTGA  CGCCGTCTTG  GGGCTGGTCT
951 GCATGTGCT  GCTGTGGTG  CTGAAGCTGA  TCGGGGACCA  CGTGCCTCCC
1001 GTCCACCCCG  AGATGCCCC  TGGTGTGCGG  CTCAGCCGTG  GGCTGGTCTG
1051 GGCTGCCACG  ACAGCTCGCA  ACGCCCTGGT  GGTCTCCTTC  GCAGCCCTGG
1101 TTGCGTACTC  CTTGAGGTG  ACTGGATACC  AGCCTTTCAT  CCTAACAGGG
1151 GAGACAGCTG  AGGGGCTCCC  TCCAGTCCGG  ATCCCGCCCT  TCTCAGTGAC
1201 CACAGCCAAC  GGGACGATCT  CCTCACCGA  GATGGTGACG  GACATGGGAG
1251 CCGGGCTGGC  CGTGGTGCCC  CTGATGGGCC  TCCTGGAGAG  CATTGCGGTG
1301 GCCAAAGCCT  TCGCATCTCA  GAATAATTAC  CGCATCGATG  CCAACCAGGA
1351 GCTGCTGGCC  ATCGGTCTCA  CCAACATGTT  GGGCTCCCTC  GTCTCCTCCT
1401 ACCCGGTAC  AGGCAGCTTT  GGACGGACAG  CCGTGAACGC  TCAGTCGGGG
1451 GTGTGCACCC  CGGCGGGGG  CCTGGTGACG  GGAGTGCTGG  TGCTGCTGTC
1501 TCTGGACTAC  CTGACCTCAC  TGTTCTACTA  CATCCCCAAG  TCTGCCCTGG
1551 CTGCCGTGAT  CATCATGGCC  GTGGCCCGC  GTTTCGACAC  CAAGATCTTC
1601 AGGACGCTCT  GCGGTGTTAA  GAGGCTGGAC  CTGCTGCCCC  TGTGGGTGAC
1651 CTTCTGCTG  TGCTTCTGGG  AGGTGACGTA  CGGCATCCTG  GCGGGGGCCC
1701 TGGTGTCTCT  GCTCATGCTC  CTGCACTCTG  CAGCCAGGCC  TGAGACCAAG
1751 GTGTGAGAGG  GGCCGTTCT  GGTCTGACG  CCGGCCAGCG  GCCTGTCTCT
1801 CCCTGCCATG  GAGGCTCTGC  GGGAGGAGAT  CCTAAGCCGG  GCCCTGGAAG
1851 TGTCCCCGCC  ACGCTGCCTG  GTCCTGGAGT  GCACCATGT  CTGCAGCATC
1901 GACTACACTG  TGGTGTGGG  ACTCGGCGAG  CTCTCCAGG  ACTTCCAGAA
1951 GCAGGGCGTC  GCCCTGGCCT  TTGTGGGCCT  GCAGGTCCCC  GTTCTCCGTG
2001 TCCTGCTGTC  CGCTGACCTG  AAGGGGTTCC  AGTACTTCTC  TACCCTGGAA
2051 GAAGCAGAGA  AGCACCTGAG  GCAGGAGCCA  GGGACCCAGC  CCTACAACAT
2101 CAGAGAAGAC  TCCATTCTGG  ACCAAAAGGT  TGCCCTGCTC  AAGGCATAAT
2151 GGGGCCACCC  GTGGGCATCC  ACAGTTTGCA  GGGTGTTCGG  GAAGGTTCTT
2201 GTCATGTGTA  TTGGATGCTG  GATGCCGCT  GATAGACATG  CTGGCCTGGC
2251 TGAGAAACCC  CTGAGCAGGT  AACCAGGGA  AGAGAAGGAA  GCCAGGCCTG
2301 GAGGTCCACG  GCAGTGGGAG  TGGGGCTCAC  TGGCTTCTG  TGGGATGACT
2351 GGAAATGAC  CTCGCTGCTG  TTCCCTGGCA  TGACCCTCTT  TGGAAGAGTG
2401 GTTTGGAGAG  AGCCTTCTAG  AATGACAGAC  TGTGCGAGGA  AGCAGGGGCA
2451 GGGGTTTCCA  GCCCAGGCTG  TGCGAGGCAT  CCTGGGGCTG  GCAGCACCTT
2501 CCCGCTCAC  CAGTGCCACC  TGCGGGGAG  GGACGGGGCA  GGCAGGAGTC
2551 TGGGAGGCGG  GTCCGCTCCT  CTTGTCTGCG  GCATCTGTGC  TCTCCGAGAG
2601 AAAACCAAGG  TGTGTCAAAT  GACGTCAAGT  CTCTATTATA  AAATAATTTT
2651 GTGTTTCTA  AATGGAAAAA  GTGATAGCTT  TGGTGATTTT  GTAAAAGTCA
2701 TAAATGCTTA  TTGTAATAAA  TACAGGAAAC  CACCCCTCAC  CCTGTCCACT
2751 TGGGTGATCA  TTCCAGACCC  CTCCCCAAC  ATGCATATGT  ACCTGTCCGT
2801 CAGTGTGTGG  ATGTATGTTT  ACAGTTCTAC  ATAAATGGGA  TCATTTTATA
2851 CATGGTGCTC  TGGAACCCAC  ATTTTTCATG  CAGTCATTG  CAGTGAATTA
2901 TTTATGTGA  TAATAAATAG  CATTAGAATA  CAAAAAATA  AAAAAAATA
2951 AAAAA (SEQ 1D NO:2)

```

## FEATURES:

5' UTR: 1-257

Start: 257

WO 02/059306

PCT/US01/42809

Stop: 2147  
3' UTR: 2150-2955

## HOMOLOGOUS PROTEINS:

Top BLAST Hits:

	Score.	E
SPLICE FORM 1:		
gi 7302719 gb AAF57797.1  (AE003802) CG5002 gene product [Droso...	418	e-116
gi 7301881 gb AAF56989.1  (AE003772) CG7912 gene product [Droso...	408	e-113
gi 5834394 gb AAD53951.1  (AF180728) sulfate transporter [Droso...	388	e-107
gi 7301216 gb AAF56347.1  (AE003749) Esp gene product [Drosophi...	387	e-106
gi 7301962 gb AAF57068.1  (AE003774) CG9702 gene product [Droso...	351	8e-96
gi 7294633 gb AAF49971.1  (AE003543) CG6928 gene product [Droso...	347	2e-94
gi 7300023 gb AAF55195.1  (AE003708) CG6125 gene product [Droso...	330	3e-89
gi 7493011 pir  T39116 probable sulfate permease - fission yeas...	267	2e-70
gi 6094367 sp O74377 SULH_SCHPO PROBABLE SULFATE PERMEASE SPBC3...	266	4e-70
gi 2626753 dbj BAA23424.1  (AB008782) sulfate transporter [Arab...	266	4e-70

## SPLICE FORM 2:

gi 7302719 gb AAF57797.1  (AE003802) CG5002 gene product [Droso...	404	e-111
gi 7301881 gb AAF56989.1  (AE003772) CG7912 gene product [Droso...	394	e-108
gi 5834394 gb AAD53951.1  (AF180728) sulfate transporter [Droso...	374	e-102
gi 7301216 gb AAF56347.1  (AE003749) Esp gene product [Drosophi...	373	e-102
gi 7301962 gb AAF57068.1  (AE003774) CG9702 gene product [Droso...	337	2e-91
gi 7294633 gb AAF49971.1  (AE003543) CG6928 gene product [Droso...	332	6e-90
gi 7300023 gb AAF55195.1  (AE003708) CG6125 gene product [Droso...	315	8e-85
gi 7493011 pir  T39116 probable sulfate permease - fission yeas...	254	2e-66
gi 9955547 emb CAC05432.1  (AL391710) sulfate transporter [Arab...	254	2e-66
gi 2626753 dbj BAA23424.1  (AB008782) sulfate transporter [Arab...	253	4e-66
gi 6502994 gb AAF14540.1 AF163975_1 (AF163975) Suta [Penicilliu...	250	3e-65
gi 6094367 sp O74377 SULH_SCHPO PROBABLE SULFATE PERMEASE SPBC3...	248	1e-64
gi 6502992 gb AAF14539.1 AF163974_1 (AF163974) sulfate permease...	244	2e-63
gi 10645530 gb AAG21641.1 AC069474_20 (AC069474) sulphate trans...	242	9e-63

BLAST to dbEST:

## SPLICE FORM 1:

gb AA582196.1 AA582196	nn44h12.s1 NCI_CGAP_GC5 Homo sapiens...	1027	0.0
gb AI690196.1 AI690196	tx33c05.x1 NCI_CGAP_Lu24 Homo sapien...	948	0.0
gb AI417381.1 AI417381	tg30c11.x1 NCI_CGAP_Brn25 Homo sapie...	902	0.0
gb AW207633.1 AW207633	UI-H-BI1-afl-c-10-0-UI.s1 NCI_CGAP_S...	890	0.0
gb AI494563.1 AI494563	qz16h03.x1 NCI_CGAP_CLL1 Homo sapien...	850	0.0
gb H15131.1 H15131	ym30b04.s1 Soares infant brain INIB Homo...	805	0.0
gb AI220943.1 AI220943	qg08c06.x1 Soares placenta_8to9weeks...	743	0.0
gb AW055140.1 AW055140	wz01c04.x1 NCI_CGAP_Brn23 Homo sapie...	718	0.0
gb AI963799.1 AI963799	wr67b09.x1 NCI_CGAP_Ut1 Homo sapiens...	718	0.0
gb N54913.1 N54913	yv34f02.s1 Soares fetal liver spleen INF...	710	0.0
gb AW008673.1 AW008673	ws71c02.x1 NCI_CGAP_Brn23 Homo sapie...	688	0.0
gb AI680222.1 AI680222	tw66e04.x1 NCI_CGAP_Ut3 Homo sapiens...	638	e-180
gb AA385773.1 AA385773	EST99540 Thyroid Homo sapiens cDNA 5...	636	e-180
gb H04038.1 H04038	yj45b06.r1 Soares placenta Nb2HP Homo sa...	626	e-177
gb AI000592.1 AI000592	os63b01.s1 NCI_CGAP_Br2 Homo sapiens...	618	e-174

BLAST to CHGI:

TA_39934	NOT ASSIGNED [Homo sapiens]	1544	0.0
TA_44508	NOT ASSIGNED [Homo sapiens]	509	e-142
TA_102843	NOT ASSIGNED [Homo sapi...	484	e-135
TA_82272	AAD14 [Homo sapiens]	44	0.018
TA_71453	ataxin SCAl [Homo sapiens]	44	0.018
TA_74619	alkaline phosphatase, pl...	42	0.070
TA_52146	NOT ASSIGNED [Homo sapiens]	42	0.070
TA_237492	NOT ASSIGNED [Homo sapi...	40	0.28
TA_158839	NOT ASSIGNED [Homo sapi...	40	0.28

## EXPRESSION INFORMATION FOR MODULATORY USE:

## SPLICE FORM 1:

Library source from BLAST dbEST hits:

AA582196 534 bp mRNA EST 05-SEP-1997	germ cell tumor
AI690196 486 bp mRNA EST 16-DEC-1999	carcinoid:lung
AI417381 470 bp mRNA EST 30-MAR-1999	anaplastic oligodendroglioma: brain
AW207633 511 bp mRNA EST 02-DEC-1999	unknown
AI494563 457 bp mRNA EST 17-MAR-1999	B-cell, chronic lymphocytic leukemia
HI5131 508 bp mRNA EST 27-JUN-1995	infant brain, 73 days post natal
AI220943 375 bp mRNA EST 29-NOV-1998	two placentae: one from 8 weeks and another
from 9 weeks post conception	
AW055140 366 bp mRNA EST 09-MAR-2000	glioblastoma (pooled): brain
AI963799 377 bp mRNA EST 08-MAR-2000	well-differentiated endometrial
adenocarcinoma, 7 pooled tumors: uterus	
N54913 486 bp mRNA EST 28-JAN-1997	fetal liver and spleen
AW008673 369 bp mRNA EST 10-SEP-1999	glioblastoma (pooled): brain

Library source from cDNA retrieval:

Human Fetal Brain  
Human Brain  
Human Pituitary Gland  
Human Heart  
Human Leukocytes  
Human Kidney  
Human Hela Cell  
Human Liver  
Human Thyroid  
Human Lung  
Human Placenta  
Human Skeletal Muscle  
Human Fetal Kidney  
Human Small Intestine  
Human Prostate  
Human Testis  
Human Adrenal Gland  
Human Bone Marrow  
Human Pancreas



## SPLICE FORM 1:

```

1  MPSSVTALGQ ARSSGPGMAP SACCSPAAL QRRLPILAWL PSYSLQWLKM
51 DFVAGLSVGL TAIPQALAYA EVAGLPPQYG LYSAFMGCFV YFFLGTSRDV
101 TLGPTAIMSL LVSFYTFHEP AYAVLLAFLS GCIQLAMGVL RLGFLLDFIS
151 YPVIKGETSA AAVTIGFGQI KNLLGLQNIP RPFFLQVYHT FLRIAETRVG
201 DAVLGLVCM LLLVLKLMRD HVPPVHPMP PGVRLSRGLV WAATTARNAL
251 VVSFAALVAY SFEVTGYQPF ILTGETAEG LPPVRIPPFSV TTANGTISFT
301 EMVQDMGAGL AVVPLMGLLE SIIVAKAFAS QNNYRIDANQ ELLAIGLTNM
351 LGSVLSSYPV TGSFGRTAVN AQSGVCTPAG GLVTGVVLVLL SLDYLTSLFY
401 YIPKSALAAV IIMAVAPLFD TKIFRTLWRV KRLDLLPLCV TFLLCFWEVQ
451 YGILAGALVS LLMLLHSAAR PETKVSEGPV LVLQPASGLS FPAMEALREE
501 ILSRALEVSP PRCLVLECTH VCSIDYTVVL GLGELLQDFQ KQGVALLAFVG
551 LQVPVLRVLL SADLKGQYF STLEEAEKHL RQEPGTQPNY IREDSILDQK
601 VALLKA (SEQ ID NO:3)

```

## SPLICE FORM 2:

```

1  MPSSVTALGQ ARSSGPGMAP SACCSPAAL QRRLPILAWL PSYSLQWLKM
51 DFVAGLSVGL TAIPQALAYA EVAGLPPQYG LYSAFMGCFV YFFLGTSRDV
101 TLGPTAIMSL LVSFYTFHEP AYAVLLAFLS GCIQLAMGVL RLAHISPHPL
151 GLGGAGTSSM SPLGWPGFLL DFISYPVIKG FTSAAVTIG FGQIKNLLGL
201 QNIPRPFFLQ VYHTFLRIAE TRVGDAVLGL VCMLLLLVLL LMRDHVPPVH
251 PEMPPGVRLS RGLVWAATTA RNALVVSFAA LVAYSFEVTG YQPFILTGET
301 AEGLPVVRIP PFSVTTANGT ISFTEMVQDM GAGLAVVPLM GLLESIAVAK
351 AFASQNNYRI DANQELLAIG LTNMLGSLVS SYPVTGSFGR TAVNAQSGVC
401 TPAGGLVTGV LVLLSLDYLT SLFYIIPKSA LAAVIIMAVA PLFDTKIFRT
451 LWRVKRLDLL PLWVTFLLCF WEVQYGILAG ALVSLMLLLH SAARPETKVS
501 EGPVLVLQPA SGLSFPAMEA LREEILSRAL EVSPPRCLVL ECTHVCSIDY
551 TVVLGLGELL QDFQKQGVALL AFVGLQVPVL RVLLSADLKG FQYFSTLEEA
601 EKHLRQEPGT QPYNIREDSI LDQKVALLKA (SEQ ID NO:4)

```

## FEATURES:

Functional domains and key regions:

## SPLICE FORM 1:

[1] PDOC00001 PS00001 ASN\_GLYCOSYLATION  
N-glycosylation site  
294-297 NGTI

[2] PDOC00005 PS00005 PKC\_PHOSPHO\_SITE  
Protein kinase C phosphorylation site  
Number of matches: 2  
1 96-98 TSR  
2 245-247 TAR

[3] PDOC00006 PS00006 CK2\_PHOSPHO\_SITE  
Casein kinase II phosphorylation site  
Number of matches: 6  
1 96-99 TSRD  
2 116-119 TFHE  
3 298-301 SFTE  
4 571-574 STLE  
5 572-575 TLEE  
6 595-598 SILD

## [4] PDOC00008 PS00008 MYRISTYL

N-myristoylation site

Number of matches: 15

1	9-14	GQARSS
2	17-22	GMAPSA
3	55-60	GLSVGL
4	80-85	GLYSAF
5	168-173	GQIKNL
6	232-237	GVRLSR
7	238-243	GLVWAA
8	307-312	GAGLAV
9	317-322	GLLESI
10	352-357	GSLVSS
11	380-385	GGLVTG
12	381-386	GLVTGV
13	452-457	GILAGA
14	456-461	GALVSL
15	543-548	GVALAF

## [5] PDOC00370 PS01130 SULFATE TRANSP

Sulfate transporters signature

77-98 PQYGLYSAFMGCFVYFFLGTSR

Membrane spanning structure and domains:SPLICE FORM 1:

Helix	Begin	End	Score	Certainty
1	11	31	0.878	Putative
2	50	70	1.193	Certain
3	77	97	1.866	Certain
4	120	140	1.767	Certain
5	148	168	1.097	Certain
6	199	219	1.547	Certain
7	247	267	1.151	Certain
8	283	303	0.683	Putative
9	306	326	1.282	Certain
10	342	362	1.277	Certain
11	372	392	1.695	Certain
12	401	421	1.366	Certain
13	449	469	1.784	Certain
14	477	497	0.915	Putative
15	542	562	0.834	Putative

## BLAST Alignment to Top Hit:

## SPICE FORM 1:

gi|7302719|gb|AAF57797.1| (AE003802) CG5002 gene product [Droso... 418 e-116

Query: 25 CSPALQRRPILAWLPSYSLOWLKMDFVAGLSVGLTAIPQALAYAEVAGLPPQYGLYSA 84  
C P+ + + PIL WLP Y L+++ DF+AG +VGLT IPQA+AY VAGL PQYGLYSA  
Sbjct: 28 CRPSTVTNKFILKWLPRYRLEYIMQDFIAGFTVGLTTIPQAIAYGVVAGLEPQYGLYSA 87

Query: 85 FMGCFVYFFLGTSRDVTLGPTAIMSLVSYFTFHEPAYAVLLAFLSGCIQLAMGVRLGF 144  
FMGCF Y G+ +DVT+ TAIM+L+V+ Y P YAVL+ FL+GCI L +G+L +G  
Sbjct: 88 FMGCFYIVFGSCKDVTIATTAIMALMVNQYATISPDYAVLVCFLAGCIVLLGLLNMGV 147

Query: 145 LLDIFISYPVIKGFSAAAVTIGFGQIKNLLGLQNIIPRPFFLQVYHTFLRIAETRVGDAVL 204  
L+ FIS PVI GFT AAA TIG QI N++GL + + F + R+ DA+L  
Sbjct: 148 LVRFISIPVITGFTMAAATTIGSAQINNIVGLTSPSNDLLPAWKNFFTHLTSIRLWDALL 207

Query: 205 GLVCMILLLLVLKLMRDHVPVHPPEMPGVRLSRGLVWA-ATTARNALVVSFAALVAYSFE 263  
G+ ++ LL++ ++D ++ + W +RNAL V F +AY  
Sbjct: 208 GVSSLVFLLLMTRVKD-----IKWGNRIFWKYLGLSRNALAVIFGTFLLAYILS 255

Query: 264 VTGYQPFILTGETAEGLPVRIPPFSVTTANGTISFTEMVQDMGAGLAVVPLMGLLESIA 323  
G QPF +TG G+PP R+PPFS T +SF EM+ +GA L +PL+ +LE +A  
Sbjct: 256 RDGNQPFRTGNITAGVPPFRLPFFSTTVDGEYVSFGEMISTVGASLGSIPLISILEIVA 315

Query: 324 VAKAFASQNNYRIDANQELLAIGLTNMLGSLVSSYPVTGSEFGRATAVNAQSGVCTPAGGLV 383  
++KAF+ +DA+QE++A+G+ N++GS V S PVTGSF RTAVN SGV TP GG V  
Sbjct: 316 ISKAFS--KGKIVDASQEMVALGMCNIMGSFVLSMPVTGSEFTRTAVNNASGVKTPLGGAV 373

Query: 384 TGVLVLLSLDYLTSLFYIIPKSALAAVIIMAVAPLFDTKIFRTLWRVKRLDLLPLCVTFEL 443  
TG LVL++L +LT FY+IPK LAA+II A+ L + + +W+ K+ DL P VT L  
Sbjct: 374 TGALVLMALAFLTQTFYFIPKCTLAIIIAAMISLVELHKIKDMWKS KKKDLFFPVVTVL 433

Query: 444 LC-FWEVQYGIAGALVSLMLLHSAARPET-----KVSEGPVLVLQFASGLSFPAMEAL 497  
C FW ++YGIL G +++ +L+S+ARP K++ V V+ L + + E L  
Sbjct: 434 TCMFWSLEYGILCGIGANMVYIYSSARPHVDIKLEKINGHEVSVVDVKQKLDYASAEYL 493

Query: 498 REEILSRAL--EVSPRCVLECHVCSIDYTVVLGLGELLQDFQKQGVALLAFVGLQVPV 555  
+E+++ R L + + +V++ + SIDYTV + + + D + A+ +  
Sbjct: 494 KEKVV-RFLNNQNGETQLVVIKGEINSIDYTVAMNIVSMKGDLEALNCAMICWNWNIA 552

Query: 556 LRVL---LSADLKG-FQYFSTLEE 575  
V+ L+ DL+ F++ +LEE  
Sbjct: 553 AGVVCRLNNDLRPIFKFDLSLEE 575 (SEQ ID NO :6)

## SPICE FORM 2:

>gi|7302719|gb|AAF57797.1| (AE003802) CG5002 gene product

[Drosophila melanogaster]

Length = 595

Score = 404 bits (1026), Expect = e-111

Identities = 233/587 (39%), Positives = 340/587 (57%), Gaps = 51/587 (8%)

Query: 25 CSPALQRRPILAWLPSYSLOWLKMDFVAGLSVGLTAIPQALAYAEVAGLPPQYGLYSA 84  
C P+ + + PIL WLP Y L+++ DF+AG +VGLT IPQA+AY VAGL PQYGLYSA  
Sbjct: 28 CRPSTVTNKFILKWLPRYRLEYIMQDFIAGFTVGLTTIPQAIAYGVVAGLEPQYGLYSA 87

Query: 85 FMGCFVYFFLGTSRDVTLGPTAIMSLVSYFTFHEPAYAVLLAFLSGCIQLAMGVRLAH 144  
FMGCF Y G+ +DVT+ TAIM+L+V+ Y P YAVL+ FL+GCI L +G+L +  
Sbjct: 88 FMGCFYIVFGSCKDVTIATTAIMALMVNQYATISPDYAVLVCFLAGCIVLLGLLNM-- 145

Query: 145 ISPHPLGLGGAGTSSMSPLGWPGFLLDFISYPVIKGFSAAAVTIGFGQIKNLLGLQNIIP 204  
G L+ FIS PVI GFT AAA TIG QI N++GL +  
Sbjct: 146 -----GVLVRFISIPVITGFTMAAATTIGSAQINNIVGLTSPS 183

Query: 205 RPFFLQVYHTFLRIAETRVGDAVLGLVCMILLLLVLKLMRDHVPVHPPEMPGVRLSRGLV 264

+ F + R+ DA+LG+ ++ LL++ ++D ++ +  
 Sbjct: 184 NDLLPAWKNFFTHLTSIRLWDALLGVSSLVFLLLMTRVKD-----IKWGNRIF 231  
 Query: 265 WA-ATTARNALVVSFAALVAYSFEVTGYQPFILTGETAEGLPVRIIPFVSVTANGTISF 323  
 W +RNAL V F +AY G QPF +TG G+PP R+PPFS T +SF  
 Sbjct: 232 WKYLGLSRNALAVIFGTFLAYILSRDGNQPFRTVGNITAGVPPFRLPPFSTTVDGEYVSF 291  
 Query: 324 TEMVQDMGAGLAVVPLMGLLESIAVAKAFASQNNYRIDANQELLAIGLTNMLGSLVSSYP 383  
 EM+ +GA L +PL+ +LE +A++KAF+ +DA+QE++A+G+ N++GS V S P  
 Sbjct: 292 GEMISTVGASLGSIPILISILEIVAISKAFS--KGKIVDASQEMVALGMCNIMGSFVLSMP 349  
 Query: 384 VTGSFGRNAVNAQSGVCTPAGGLVTGVVLVLSLDYLTSLFYYPKSAALAAVIIMAVAPLF 443  
 VTGSF RTAVN SGV TP GG VTG LVL++L +LT FY+IPK LAA+II A+ L  
 Sbjct: 350 VTGSFTRTAVNNASGVKTPGGAVTGALVIMALAFLTQTFYFIPKCTLAIIIAAMISLV 409  
 Query: 444 DTKIFRTLWRVKRLDLLPLWVTFLLC-FWEVQYIGILAGALVSLMLLHSAARPET----- 497  
 + + +W+ K+ DL P VT L C FW ++YGIL G +++ +L+S+ARP  
 Sbjct: 410 ELHKIKDMWKSKKKDLFPFVTVLTCMFWSLEYGILCGIGANMVYILYSSARPHVDIKLE 469  
 Query: 498 KVSEGPVLVLQPASGLSFFAMEALREEILSRAL--EVSPPRCLVLECTHVCSIDYTVVLG 555  
 K++ V V+ L + + E L+E+++ R L + + +V++ + SIDYTV +  
 Sbjct: 470 KINGHEVSVDVKQKLDYASAEYLKEKVV-RFLNNQNGETQLVVIKGEINSIDYTVAMN 528  
 Query: 556 LGELLQDFQKQGVAFVGLQVPVLRVL--LSADLKG-FQYFSTLEE 599  
 + + D + A+ + V+ L+ DL+ F++ +LEE  
 Sbjct: 529 IVSMKGDLEALNCAMICWNWNINIASAGVVCRLNNDLRPIFKFDLSLEE 575 (SEQ ID NO :7)

Multiple sequence alignment of both splice forms and the top blast hit:

Splice Form 1 ~~~~~~MPSSVTAL GQARSSGPGM APSACCCSPA ALQRRLPILA  
 Splice Form 2 ~~~~~~MPSSVTAL GQARSSGPGM APSACCCSPA ALQRRLPILA  
 gi7302719\_pe MRAEDNLYR EQLPNVSTLI ...RDGGRKL .....CRPS TVTNKFPILK

51 100  
 Splice Form 1 WLPSYSLQWL KMDFVAGLSV GLTAIPQALA YAEVAGLPPQ YGLYSAFMGC  
 Splice Form 2 WLPSYSLQWL KMDFVAGLSV GLTAIPQALA YAEVAGLPPQ YGLYSAFMGC  
 gi7302719\_pe WLPRYRLEYI MQDFIAGFTV GLTTIPQAIA YGVVAGLEPQ YGLYSAFMGC

101 150  
 Splice Form 1 FVYFFLGTSR DVTLGPTAIM SLLVSFYTFH EPAYAVLLAF LSGCIQLAMG  
 Splice Form 2 FVYFFLGTSR DVTLGPTAIM SLLVSFYTFH EPAYAVLLAF LSGCIQLAMG  
 gi7302719\_pe FTYIVFGSCK DVTIATTAIM ALMVNQYATI SPDYAVLVCF LAGCIVLLLG

151 200  
 Splice Form 1 VLRL.....GF LLDFISYPVI KGFTSAAAVT  
 Splice Form 2 VLRLAHISPH PLGLGGAGTS SMSPLGWPGF LLDFISYPVI KGFTSAAAVT  
 gi7302719\_pe LLNM.....GV LVRFISIPVI TGFTMAAATT

201 250  
 Splice Form 1 IGFGQIKNLL GLQNIIPRPF LQVYHTFL.R IAETRVGDAV LGLVCMLELL  
 Splice Form 2 IGFGQIKNLL GLQNIIPRPF LQVYHTFL.R IAETRVGDAV LGLVCMLELL  
 gi7302719\_pe IGSAQINNIV GLTS.PSNDL LPAWKNFFTH LTSIRLWDAL LGVSSLVFLL

251 300  
 Splice Form 1 VLKLMRDHVP PVHPEMPGV RLSRGLVWA. ATTARNALVV SFAALVAYSF  
 Splice Form 2 VLKLMRDHVP PVHPEMPGV RLSRGLVWA. ATTARNALVV SFAALVAYSF  
 gi7302719\_pe LMTRVKD... .....I KWGNRIFWKY LGLSRNALAV IFGTFLAYIL

```

301                                     350
Splice Form 1 EVTGYQPFIL TGETAEG LPP VRIPPF SVTT ANGTISFTEM VQDMGAGLAV
Splice Form 2 EVTGYQPFIL TGETAEG LPP VRIPPF SVTT ANGTISFTEM VQDMGAGLAV
gi7302719_pe SRDGNQPF RV TGNITAGVPP FRLPPFSTTV DGEYVSFGEM ISTVGASLGS

351                                     400
Splice Form 1 VPLMGLLESI AVAKAFASQN NYRIDANQEL LAIGLTNMLG SLVSSYPVTG
Splice Form 2 VPLMGLLESI AVAKAFASQN NYRIDANQEL LAIGLTNMLG SLVSSYPVTG
gi7302719_pe IPLISILEIV AISKAFS..K GKIVDASQEM VALGMCNIMG SFVLSMPVTG

401                                     450
Splice Form 1 SFGRTAVNAQ SGVCTPAGGL VTGVLVLLSL DYLTSLFYI PKSALAAVII
Splice Form 2 SFGRTAVNAQ SGVCTPAGGL VTGVLVLLSL DYLTSLFYI PKSALAAVII
gi7302719_pe SFTRTAVNNA SGVKTPLGGA VTGALVLMAL AFLTQTFYFI PKCTLAAIII

451                                     500
Splice Form 1 MAVAPLFDTK IFRTLWRVKR LDLLPLCVTF LLC.FWEVQY GILAGALVSL
Splice Form 2 MAVAPLFDTK IFRTLWRVKR LDLLPLWVTF LLC.FWEVQY GILAGALVSL
gi7302719_pe AAMISLVELH KIKDMWKSCK KDLFPFVVTV LTCMFWSLEY GILCGIGANM

501                                     550
Splice Form 1 LMLLHSAARP ET....KVS EGPVLVLQPA SGLSFPAMEA LREEILSRAL
Splice Form 2 LMLLHSAARP ET....KVS EGPVLVLQPA SGLSFPAMEA LREEILSRAL
gi7302719_pe VYIYSSARP HVDIKLEKIN GHEVSVVDVK QKLDYASAAY LKEKVV.RFL

551                                     600
Splice Form 1 ..EVSPPRCL VLECTHVCSI DYTIVVLGLGE LLQDFQKQGV ALAFVGLQVP
Splice Form 2 ..EVSPPRCL VLECTHVCSI DYTIVVLGLGE LLQDFQKQGV ALAFVGLQVP
gi7302719_pe NNQNGETQLV VIKGEEINSI DYTIVAMNIVS MKGDLEALNC AMICWNWNIA

601                                     650
Splice Form 1 VLRVL..LSA DLKG.FQYFS TLEEA EKHLR QEPGTQPYNI REDSILDQKV
Splice Form 2 VLRVL..LSA DLKG.FQYFS TLEEA EKHLR QEPGTQPYNI REDSILDQKV
gi7302719_pe SAGVVCRLNN DLRPIFKFDL SLEEVVAGHF DSPSNTASTV TIEA~~~~~

651
Splice Form 1 ALLKA
Splice Form 2 ALLKA
gi7302719_pe ~~~~~

```

# Hammer search results (Pfam):

Model	Seq-from	Seq-to	HMM-from	HMM-to	Score	E-value	Description
Sulfate_transp family	137	468	1	328	229.9	1.3e-66	Sulfate transporter

1 GGTCCCCGCG GCCCTCGGCC TTGCTCGGGG CCAAGGGACC GCGGACGGTC  
 51 AGGTGGCGCA GGGTCTCCTC CGGAGACCCC AGGATCCGGA GCCAGCGGCC  
 101 TTGTGGGCAG GGGCCGGGGG CAGGGGAGTG GATTTTGCCC GGAGCGGAGC  
 151 AGGCCGGGGG CAGTGGGGGG CTGGGGGTGA GGGTGGCTGG CTCTGCGCGC  
 201 GGGCGCCGGG GCCCTGGAAG ATGCTGCGCA CCTGAATTAA CCGGGCGCCT  
 251 CTGATGTCCT CCCAGAAGCA ACTAGAACTC CAGGGCTGTG AAAGCCACAG  
 301 GTGGGGGCTG AGCGAGGCTT GGCTCAGGA GCGGAGGACC CCCCCCCCCC  
 351 CCCCCCTCGAG CGCCGCGATC CACCGTAGCG GGTGGAGCCC GCCTTGGTGC  
 401 GCAGTTGGAA AACCTCGGAA GCGCCGCTGG ATCTCCTGGC TGCCACCCGC  
 451 ACCCCCCGCG AGCTACGGTG CGCCCGCGGG CCCAGCTTCT CTCTGCGCTG  
 501 CTCCCCGTTA AATTCCCTGG GGAGACGGAA AAAAAGGCAA AGGAAGTCGG  
 551 TTCTCCAGGG GCCAGAAGTG TTGAGCCTAA TTAGTCTTCA GACTTCTCAA  
 601 TGAGGAATCG CTTATCAGTT TCTTATCTGG GAGAGTTGAG GATGGAGGGA  
 651 CAGAAGGCAC CCAGGATTG CACGGGGGGG GATTACAGGA GAGAGGGTGA  
 701 TGAGGGACGG GGTGGGCCTT CCAGTCTTGG CCCAGTCCCC ATCTGCACA  
 751 CATTGTTGGC TTCTCTTAG AGCCGTTTCG CCCCCTGGGG AGGGGAGACC  
 801 CATAGTGACC TCTCTGACA CCCGCCGACC CTGACCAGTG TTGCCGGGT  
 851 CTTCAAAGGC CACGCTCTGA CTGCTGGTCT GTGTACCTG CACCCCCAG  
 901 CCCCACCGTA GAGATGCCTT CTTCCGTGAC GCGCTGGGT CAGGCCAGGT  
 951 CCTCTGGCCC CGGGATGGCC CCGAGCGCCT GCTGCTGCTC CCCTGCGGCC  
 1001 CTGCAGAGGA GGCTGCCCAT CCTGGCGTGG CTGCCCAGCT ACTCCCTGCA  
 1051 GTGGCTGAAG ATGGATTTCG TCGCCGCGCT CTCAGTTGGC CTCACTGCCA  
 1101 TTCCCCAGGC GCTGGCCTAT GCTGAAGTGG CTGGACTCCC GCCCCAGGTG  
 1151 AGGCGTCTGA CCCTGCTGCC AGCCATATCT CAGAAACAGT GCAGAATACA  
 1201 CAGTATCAAT CCCAGACACC ATCAGCGATT CCAGGTTTCC AGCCCCTGGG  
 1251 CCCCAGGAA CCTTTGGTTT ACAGTGTGTG ACGCAGATTG TCTCTGGGCC  
 1301 GACCCAGGCT CCTATGCTG TTTGGTACAC ACAGACACTG AGCTGGTTAT  
 1351 GGAGGGGCCA GCGAGATGAC TCATGGAGGC CTCAGGAGTT CAAGACCAGC  
 1401 CCGACCAAAA TGGTGAAACC CCGTCTCCAC TAAAAATACA AAAATTAGGC  
 1451 TGGGTGCGGT GGCTCAAGCC TGTAATCCCA GCACTTTGGG AGGCCGAGGC  
 1501 AGGCGGATCG CAAGTCCAGG AGATGGAGAC CATCCTGGCT AACAGGGTGA  
 1551 ACCCCGCTCT CTAATAAAAA TACAAAAAAT TAGCCAGGTG TGGTGGCGGG  
 1601 TGCTGTAGT CCCAGCTACC TGGGAGGCTG AGGCAGGAGA ATGGCGTGAA  
 1651 CCGGGGAGGC GGAGCTTGCA GTGAGCCGAG ACTGTGCCAC TGCCCTCCAG  
 1701 CCTGGGCGAC AGAGCGAGAC TCCATCTCAA AAAAAAAAAA AAAAGAAATGC  
 1751 TTCTCAGAC TTGGACACAG CACACGGGCC TGACCGACC CCTCTGCCTG  
 1801 GCTGTCTGCA CCCTGAGGCC CCAGTTGAGT GCTGCTAAAA AAGTGGCCTC  
 1851 CTGATCACTG CAGTCCACC CACAGGGCAG GCGGGTGCAC CTTTAACCTG  
 1901 GGCTGGACA CAGTGACAC CCACACATCC CGAGCTTGA CACGCACACT  
 1951 AGGGAGCTGG TGGATGGGCC TCGGCCTCCT GAGTGTCTAC CACCCTCTCT  
 2001 CCCCACAGTA TGGCTCTAC TCTGCCTTCA TGGGCTGCTT CGTGTATTTC  
 2051 TTCTTGGGCA CCTCCCGGGA TGTGACTCTG GGCCCCACCG CCATTATGTC  
 2101 CCTCTGGTCT TCCTTCTACA CCTCCATGA GCGCGCTAC GCTGTGCTGC  
 2151 TGGCCTTCCT GTCCGGCTGC ATCCAGCTGG CCATGGGGGT CCTGCGTTTG  
 2201 GGTGAGGCTC TACCTTCTTG CCAAGGGGAT GCCCTCGACC TCAGCATTTG  
 2251 CTTGTTTGCA TTCAAGTCT ATCCCCGTGT GCGTGTGTGT GCGTGTGTG  
 2301 GGGGTGTGGG TATGTATGTG TGTGTGTGTA GGTGGGTGGG TGGTGGAGGG  
 2351 GGTGGGGCAC TTGGCTCCTT AGTCTACTAT TTTACTGATT AGAGGCCAGG  
 2401 ACATTGGAGA AAGTGACCTG TGGCTCAGAC CCCATATGCC CCNNNNNNNN  
 2451 NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNTTCATTTC  
 2501 CTACCCCGTC ATTAAAGCTT CACCTCTGCT GCTGCCGTCA CCATCGGCTT  
 2551 TGGACAGATC AAGGTAGGCA CGGCGCCAC CCAGGGCACT GCTCTTTGGC  
 2601 CACTGCTCGT TGGCACAGGG ATGGCGGGAG CAGGACTGAG GCCAGTCCTG  
 2651 ATCCCTGTGG CCAGTGGACG TCTTGCTGTT TCAGATTGTC TTCCATGGGT  
 2701 CAAGAAGCAC GCGGTGCTCT CATGGGTCCC CTGTAAATAA AATGACCCTC  
 2751 CTGGGAGGGA TGTCACGTGA TGGTTGGATT TCACAGCGGG TAACTTGGGG  
 2801 GCCGGTAATT CCATCCCCCT GCTCTTGCCC GAGTTTCCGT GCCAGTGTGC  
 2851 TTGGCTGGCT CTGTGACGTG GCTCTGTTCT CCCTGCACTG GGCACACCCA  
 2901 GCAGGCCCCA CCAGTCATGA GCATGCTGCT AGAATTTCTA TAGGCAAATT  
 2951 ATTTCCCAT GCCAATTTAG TTAGATGGTT TTGTTTTGTG CTTTTTTTTT  
 3001 TGGTTTCTGT TTAATATTTT TTAATGCCA TGCTTTTAT ATTTTCTTTT  
 3051 AAAACATTTA TAATAAGTAC TAAATCAGT CATCAGCTGA GGGTGAATT  
 3101 TATTCTGTTT TTGCTGGGTT GTAAGTTCTT CGAGGGTAGG AACTGTCCTT  
 3151 CTCCAGCCA TGGCCGCCCA GCATTGGGCT GTGTCAGTAG GGGGGTGTCT  
 3201 AGTGGGGTGT GTGTGGAGTG AAGTGAGCCC AGTCCACAG ATGGGACCAT  
 3251 GCGGCCCTCA TGGCAGACTA GGGTCACATG CTGCCTCTG ACCCTGTGTC

3301 ACTGCAGGTC ATCTTCCCCA AGCCAGGCCC TGTTCAGGC TGGCCTGAGA  
 3351 CAGTCTTCCC TGATGGAGGT ACCATGAGAA GACCAAGGAC AGGAGAGTGT  
 3401 GTGTGAGAGT GTGTATGAGT GTGTGTGTGT GTGAATGAGT GAGTGTGTGA  
 3451 GAGTGTGAGT GGGTTTGAGG GAGTGTGTGT GTGCGTGTGT GAGTGTGTGT  
 3501 GAGTGTGGGT GTGAGTGAGT GCGTGTGTGT GAGAGAGTGT GTGAGTGTGT  
 3551 ATGAGTGAGA GTGTGAGAGT GGGTGTGGGT GTTAGGGAGT GTGTGAGTGT  
 3601 GCGCGCGCGT GTGTGAGTGT ATGAGTGTGA GAGTGTGTGT GAGTGTGTGA  
 3651 GAGTGGGTGT GGGTGTGAGT GTGCGTGTGA GTGTGAGAGT GTGAGTGTGT  
 3701 GAGTGTGAGA ATAAAGTAGA CACTTTTTCG ACTCTTGCTA CGTGCAGAGC  
 3751 ACTGGGCGAGG AACTCTCTCA TGTGTAATTC TCAACACACC CCGGGAGGTA  
 3801 GATTTATCAT TATTTCCATT TGAGAGAGGA GGGACCAACT TAGGTGTGGG  
 3851 TGAGCGTGAT TTGTACGTTA TCTACACGCA TCTCTCAGGG TTAGCTTGGC  
 3901 AAATGCTGTT TCAGGGCATG GTTGGTTCTT TAATCTGGAA ACATCATTTT  
 3951 TGGTGTCAAG AATGTTCTTT TGTAGGATCC CAGTGTGAGT GGAGAGCGGA  
 4001 GAGTGGAGAG TGAAGGCGAT CCCTTGTTCA TGCCTTCATA CTTGGCAACC  
 4051 CTAGCCCCGC CCAGGGACTC TGCAGCCATC TGGGGGGAGG GCGTCTCTCC  
 4101 TGACAGGCCC AGGACAGAAG ACCCTACCCC AACAGTCCCA GAGTCCGCCC  
 4151 CCCAGGATGT CCTAACCCCA CCCACCCTGT ACGCAGCAGT TTTAGGGCAG  
 4201 GGTCTTGTAG CCTGTGGCCA TGGGATCCAG GGCCTCAACT TTCCCTCTG  
 4251 ACTCCTGTTG GGTCTCAGGC GATGTAAAGA ACTAAAGTGC AAGCTGTGCC  
 4301 TGCACTGTGG ATCCAGCTA CGTGGGAGGC CGAGGTGGGA GGATCACTGG  
 4351 AGCCAGGAG TCGAATCCCA GCCTGGGCCA CAGAGCAAGC AAGACCCCA  
 4401 CCCCCAACCC CCGGCGCGCT CTCTAAAGAA AAAAAACTA AACTAAACA  
 4451 CAGGATAGAG TGCTTCTCC TGCCAGGAC CTCAGAGCTG GTATCGTGGT  
 4501 GGGAGGCTCC TACTTTGCCG AGGATTCCCC AAGCTGGTTT CTTGAAGCCC  
 4551 CTCAGAGCCC TCCACATCTG CACACTACGA AGAGATTTT CCTCCCGCAG  
 4601 CAGCGGAGCT GGGGGGTGGC GGGGCACCTA GTGAGGGAGA CATTCTCAAG  
 4651 CCAATGGCAG CAAGGGTCTA CACTGCAGGG GGCCTGGCTG CTGAGTGGCC  
 4701 TGGAGCCAAT GGGGGTGGGG CCAGGCAGCC CATGGCCCTT GGCCATCAAG  
 4751 CTGTAGAACC TGCTGCCTG CTGTGGGCTG ACCTACATTG TTTTATGA  
 4801 GAAAAGTAAT TTAGAGAAAA ACATATCACT GACCCAGTAA TTGTGAAGTA  
 4851 CTGTCTCCCA CGAGGGTAGC TTTGATCTCC TGCCCTAGGG GCGTTGGGA  
 4901 GTGGGCAGGT GGACCCCTGA GCCCTAAGA TGGGCCCCGG GAATAAGGGT  
 4951 TGGGAGCAGG GCCGGGGGAC ACTGTCTCAG CCCTAGGGGA GGTGGGCGGG  
 5001 GAGCTGGGGA CAGATGGCCT TGGTTGGGA GCATAGCCTC TGATCAGCAT  
 5051 CTCTGTGTTT GGACAGAACC TGCTGGGACT ACAGAACATC CCCAGGCCGT  
 5101 TCTTCTTGCA GGTGTACCAC ACCTTCCTCA GGATTGCAGA GACCAGGTAC  
 5151 CCGGGGCTTT GTTCTCCTT CCTATAAGGA AGCTCCTTCT TCCACACCTC  
 5201 CTCTCCCGGC CCCCACCTCA GTTCCCCAC CCCTGGTGAC TGCTCAAACA  
 5251 GGGGTCCCCA GAGCAGCCCC ATCAGCAGCA CCTGCAAGCT GGCAAGAAAT  
 5301 GAGACTCTCA GTCCTTTGCA GACCTGCCCT ATCGGAGCCA GAGGTTAAC  
 5351 AGGAGGCCGG TGGTCTGCGC GCAGGGGAGA GTTCAAGAAG CTCTGCTCTG  
 5401 GGGAGCTGGA GCAGGGCAAT GTTCTCTCT CCACAAAGCT TTCTTGGGG  
 5451 ACAGGAATGG GGAGAGTCTG TGTAAAGAAA AAGAAGGCAG TGTGTCTCCC  
 5501 CCAGGCTGTG ATTTGTTAAG GAGGAGAACA CAGGGCGTGG GGAGCTAACC  
 5551 CAGACAGAAC GCAGGTGGTG CAGGGACGGC AGGTGGAGCT GTGATGAGAG  
 5601 ATGAGGAGAC CCAAGCACTT CCCTCAGGTG GTTTAAAGAG CCTCTGATGG  
 5651 CCGGACACTG TGGCTCACAC CTGTAATCCC AGCACTTTGG GAAGCCAGG  
 5701 TGGGCAGATC ATCTTAGGTC AGGAGTTCAA GACCAGCCTG GTCAACATGG  
 5751 TGAAACTCCG TCTCTACTAA AAATACAAAA ATTAGTCGGG TGTGATGGTG  
 5801 CGTGCCTGTA GTCCAGCTG TTCAGGAGGC TGAGGCAGGA GGATTGCTTG  
 5851 AACCTGTGAG GCGGAGGTTG CAGTGTGCTG AGATCACGCC ACTGCACCTC  
 5901 AGCCCGGGTG ATAGAGTGAG ACTCCATCTC GCCTCTGCCC TGCCCCCGG  
 5951 AAAAGAAAAG CCTCTGATGA GGGGTACCTC CCTGCCAGAC CATCCAGCGG  
 6001 GAAGGCAGGA TGCCTCTCTA CCTCTCTGGC TGAAGGGGC TGGAGGAGGA  
 6051 AATGATCTAG GGGAGCTATA GAGATGGCTG CCCAGTGCTG TGGCCTGGAG  
 6101 GGAGTGGGGG CGATCCTGGA CCATCTTCCC CTCCCTCTG ATCTGCTGCC  
 6151 GAAGCTTCCC AGCAGCTGAA CCCAGCTGGA AGCCAGTGGG TTCTTTGCTC  
 6201 TTCAGAGGCA CCAGTGGGCA GGGGTGGGCC AGATGGGGGA GCAGTGATTA  
 6251 CGGAGCCTGA CCAGTCAGAG AGCGTCACGG GCTGTCTAGT AAAAGGGGCC  
 6301 TTAGTGACCG TTTAGTCCTA AGCCTTCTAG AATGTCTAGA GCAGGGTGTC  
 6351 CAGTCTTTTG TCTTCCCTGG GCCACACAGG AAGAATTGTC TTAAATACA  
 6401 CTAATGATAG CTGATGAGCT ACAAATAAAT CATTGAAAGA ACTCATAATG  
 6451 TTTTAAGAAA GTTTACAAAT TTGTGTTGGG CCACATTCAA AGCCATCCCA  
 6501 GGCTGCACAT GGCCACAGG CTGTGGGTTT GGCAGACTTG TTCTAGAGCC  
 6551 ACCTGCAGAG AGCATCAGCT GCTCAGGCTG GTCCCCATT CCTGTGGTCA

6601 CTCACCAGAT CCTGTTCCAT AGACTTGAGT CCCAGAGAGC TACGGGAGTG  
6651 AAAATGTGAG CACGTGCTC TGCCATCATC ATGACCCGGG CCTCTCCACT  
6701 GCCTGTCCTG CCCAAGCCTG TGTACTTACT GAATTTTGAA TTGAGTTTGT  
6751 AAGTAGAACA GGGATGCATA TTGCTTAGGA GAAAACATCT CAGCCTAGTC  
6801 TGTGTTTGGT CTTGCAGAGT AGCTTGCTAA AGTTCCCTGAG CTTTAGTTAA  
6851 GTGTCTGTGA AATGGTGAAA ATATTGAAAA TGCTTTATTT CTTGTGATA  
6901 CTCACTGTGG TCTGGGACCA GCAGCCTGGG CATTGGGCCT CCCTGGGAGC  
6951 TTCTTAGTGC CTCTCAGCTT AAGCAGCAGC AGGACGCTTG AGTCAGCAGT  
7001 AGGCCTGGGC CGAGCTGGGT GGTGACCAGT CCTCTGCCTG TCCACAGGGT  
7051 AGGTGACGCC GTCCTGGGGC TGGTCTGCAT GCTGCTGCTG CTGGTGCTGA  
7101 AGCTGATGCG GGACCACGTG CCTCCCGTCC ACCCCGAGAT GCGCCCTGGT  
7151 GTGCGGCTCA CCGTGGGCT GGTCTGGGCT GCCACGACAG GTGAGGGGCC  
7201 TCTGGCTGAC ATCGTATGCA ACCTGGCTG CAGGTGGGG TCACTGGGG  
7251 AGTCCTAGTC CCACCCTAGG GATTCTCAGC TCATTGGTCT GGGTGTCACT  
7301 TGAGCATTGG GACATTTAAA ACACCACACC AAACCTCTGG ACATGTACTT  
7351 TTTATTTAAT TAATTAATTA ATTATTTTTT GAGACAGAGT TTCGCTCTGT  
7401 CGCCAGGCT GGAGTACAGT GGTGTGATAT TGGCTCACTG CAACCTCCGC  
7451 CTCCAGGTT CATGTGATTC TCCTGCCTCA GCCTCCTGAG TAGCTGGGAT  
7501 TACAGGCGCA TGCCACCACG CTCGACTGAT TTTTGTATTT ATAGTAGAGA  
7551 CGGGGTTTCA CCATGTTGAC CATACTGGTC TCAAACCTCT GACCTCGAGT  
7601 GATCTGCCCTG CCTCGGCCTC CCGAAGTGCT GGGATTACAG GCATGAGCCA  
7651 CCGCCCGAC CCAAACCTCTA GAACATTTAA AACTCTGAGC CAAACTCGGT  
7701 GGGTTCCTGAT GCAGGGTCAG AGCTGAGAAC TGATGCATGG CCAGCTGTGG  
7751 CCGTTCCTCAA CCTGAAGCAG TTTTGCCCCC CTGCTGACAT GTGGCAATGT  
7801 TGGGAAACAT TTTTGGTGGA TACCACCGGG CGAGGCAGTT GCTGGCAACT  
7851 AGCGAGGAGG GGCCAGGAAT GCCACTAAGC CCCTGACGGC GCACAGCCTT  
7901 CCACAGCAAA CAGTGATCCG GCCCAAAATG TCGGCGGTGC CGAGGTGGAG  
7951 AAACACAGTC TACTGGCAGG TCCTGGGGAG ACAGAACAAA TTCCGGGGAG  
8001 ACCAACAGGC TCTGTTTGAA TTGTGCAGCT TTATGGGGGT ACAACTTCAG  
8051 CTTACAGGCGG GGAGGCTGCA GGCTGAGCCA CAACCAACCA GGGGTCTTT  
8101 GAGAAAGGGG TGGTGAGGGC TCGCTCGTGG AGGAGCAGGG CTGTCAGCAG  
8151 GTGAAGGACC ACCCTGGGGG AGGTGGAAGC CTCTTCTCAG TGCATCCGGG  
8201 ATGGGCATGG GTCGCCCTGGC TGAGCACCGG GCAGGGATGC CCAGAGAAGA  
8251 AGGCACACCC AGGAAAGTCA GGAAGCATAG GTGACTCCAA ACCTTCTCAG  
8301 AGCCGAGGGG ACAGATTAAG CTCAGAATAA CCCGAGCTTG GTCTTGCTGT  
8351 TTGGGGGCTG CAAAATCCAA CAGCCACAGA AGAGAGAGGG TGGCTGGATT  
8401 TAGCTCTTGA GTCACCTCTG CCTGCCCAGG CTGGTCTGTT CTCTGTGGTC  
8451 TCTCCCTGGA TCCTGTTTCA CAGACTTGAG GCCGTGGTGA AACTGAGAG  
8501 TCTGGTGGGG TGCAGAGGAT GCTGGGTTA GAGTGGAGAT GTCCAGCTTC  
8551 CAGTCCCAGC CTGGCCTGGG TTAGACTTG CAAAGTAGCT TCCAAAGTTC  
8601 CTGAGCTTTA GTTCAGTTGT CCGTGAAATA GTGAAATGC TTTGTTATCT  
8651 TTGTATCTT TTGTTTTTTG TTTTTTTTTG GAGACGGTGT CTCACCTCTG  
8701 TTGCCCAGGC TGGAGTGCAC TGGCCGTGAT CTTGGCTCA CTGCAACCTG  
8751 CCTCCTGGTT TTCAAGCAAT TATTCCTCCC TTAACCTCC CCGAGTAGCC  
8801 TGGGATTTAC AGGCCGTGCA CCCACCACAC CCGGGCTAAT TTTTGTATT  
8851 TTTGAGTTCA AGATGGGGGT TTCACCATTG TTGGCCCATG CTGGGTCTCG  
8901 AAACCTCCCN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN  
8951 NNNNNNNNCC CTAACCTGTA AAGGCAGTGT CTCCGCAGCC AGAAAGGTCT  
9001 TTTAAGATGT TTAGACATCG GTAATGTCCA GAAACTTCAG AAACATTTTA  
9051 CAGTGTAGGG CTGCTGACTC CTTTCTCCCC AGTGTGTGTA GCATTGTCTT  
9101 AGGAAACTAC TTAATAAGCT TTTGATTCTT ATAACAGCTT TGAGAATGAG  
9151 TTTATTATTT TTTCCATTTT GTCAGTGAGG AAAGTGAGCC TATGAAAGAT  
9201 GAAAAATAAA ACAGAACTTC TCAAGGCCAC ACGACTGGGA AGAGGCAGAA  
9251 CAAGGACTTC ACCCGAGTCG TCTGTCTTTT TTTCTGAATG GTCTGTCTAG  
9301 AAGTGTGTCA ATGTTATGAT CCTCTCAAAG AACCAGCTTT TGGTTTCATT  
9351 GATCTCCCTT TTGCTTTGGT TTTCTAGTTC ATTGATTCTT GCTTGGGCCT  
9401 TTATTATTTT CTTTCTTCTA TTTATTTTCA GTTAACTTTA TCCTTTTTTT  
9451 TTTTTTTTTT AAAAAAAAT TTTTTTTTTT GAGACGGAGT TTCGCTCTTG  
9501 TCCCCCAGT TGGAGTGCAA TGGTGTGATC TCAGCTCACT GCAACCTCCG  
9551 CCTCCCGGGT TCAAGTGATT CTCCTGCCTC AGCCTCCCAA GTAGCTGGGA  
9601 TTACAGGCGT ATGCCACCAT GCCTGGCTAA TTTTTGTAT TTTAATAGA  
9651 GACGGGGTTT CACCATGTTG GCCAGGCTGG TCTCGAACTC CTGAACCTCAG  
9701 GTGATCCACC CGCCTCAGCC TCCCAAAGTG CTGGGATTAC AGGCGTGAGC  
9751 CACTGTGCCC GGCCTTTTTT TTCAGATTCT TAAGTGGGAA GCCTGGGTCT  
9801 TTGATTTGAG ACCTTCTTCT TTTTCTAATA TGGTTGTTTA AATGCTGTAA  
9851 ATTTCACTCC TAGCATTACT TTAGCTGCAT CCCACAGTTT TTATATATAT



```

9901 GCTGTTCTTT AATTTTATT CGGTTCTAAA TACTTTGTGA TTCCATTTT
9951 GACTTCTCCT TTGACCTGTG GATTATTTAG AAGTGGATTA TATGATTCT
10001 TCTGTGAAGC ATGGGAGTTA TTTAGTTTCC AAATAGTTGG GGGTTTCCCA
10051 ATTAGTACTT TGCCATTGAT TTCTAATTTA GTTCCATTGT GATAAGAAAA
10101 CGTACTTAGT GATTTAAATC TGTTTTCTCT TATTGAGAGT TGTATTATGG
10151 CCCTGAATAT GGTCTATCTT AGTGAATATT CTGTGTGTGC TTGAGAACAT
10201 GTTTTCTGCC ATTATTGGGT GGAATGTTCT GTAAATGTCA GTTAGGCTCG
10251 GTTGGTTATC TCAGGCTACT ATATTCTTGC TGATTTTCTT TCTACTTGT
10301 CTGTTGATTA TTAAGAAG AATGTTGAAA TAACTTCAGC TATAACCATG
10351 ACTTTGTCTA TTTTCTTTC TTACTTACCA GTTATTGTTT CATATATTTT
10401 GAAGCTCTGT TTTTGTATG AAAACATAT AGAATTATAA TTTTCTCTTG
10451 GAATGATACT TTTTCTCTG GGCAATATTC TCTTCCCTGA AATCTACTTT
10501 GATACTAAGA GAGCTACTCC AGCTTTCATT AGTGTTAGCA TATCTTTTCC
10551 CATCTTTTAA CTTTTTTTTT TTTGAGACAG AATCTCTCTC TGTGCCCAG
10601 GCTGGAGTGC AATGGCGTGA TCTTGGCTCA CTGCAACCTC CACCTCCCAG
10651 GTTCAAGCAA TTGTTGTGCC TCAGCCTCCC AAGTAGCTGG AATTACAGGC
10701 TTGCACCACC AAGCCAGCT AATTTTGTG TTTTGTAGT CAATGAGGTT
10751 TCACCATGTT GGCCAGCCTG GTCTCAAATT CTTGGCCTCA AGTGATCCAC
10801 CTGCCTTGCC CTCCCAAAGT GCTGGGACAC CACACCCAGC CTCCTATCTT
10851 TTTACTTTTA ACGTATTTAA TTGATTGTG TCTTGTGTG TAATGGTATG
10901 TTTTATGGGC AGCATATAGA TGGGATTGT TTTAATCTC ATAATCTCTG
10951 CCTTTTAATT GGAATCTTAT GCCATTATA TTTAATGTGA TTATTGGTCT
11001 CTTTGACTTT AGATCTACCA TGTGCTGTT TGTTTCTGT TTGTCTCATT
11051 TGTTCCTTGT TCCCTTTTCC CCCTCCTCT ACCTCTACTG GATTATTTT
11101 TATGATTCCA TTTTACATCC TTTATTGGCT GATTAGCTGT AACGGCTATA
11151 TTTAGTTTTT GTTTGTTTGT TTCTAAGTGA CTGCTTTAAG GTTGTGGTA
11201 AACATCTTTA TCTTGCTACA GTATACATTA AACTTCAACT TCATGTGATA
11251 TAAGAACCTT ATAACAGGCC AGGCGCAGTG GCTCACACCT ATAATCCTAG
11301 CACTTTGGGA GGCTGAGGGA GGCAGATTGT CTGAGCTCAG GAGTCTAAGA
11351 CCAGCCTGGG CAACATGGCG AAACCCCATC TCTACCAAAA ATACAAAAAA
11401 TTAGCCAGGC ATGGTGGTGC ATGCCTGTAA TCCCAGCTAC TGGGGAGGCT
11451 GAGGTGGGAG GATTGCTTGA ACCTGGAAGG CAGAGATTGC AGTGAGCCAA
11501 GATCCGTCAC TGAAGCTGCA AACTCCTGGG CTCAAGTAAT TCTCCTGCCT
11551 CAGCCTCCTG AGTAGCTTGG TCTACAGGCA TGCAACACCA CATCTGGCTA
11601 ATTTTTTAAA TTTTTTTATA GAAACGGGCT ATGTTGCTCA GGCTGATCAA
11651 AGGTTCTTGG CCTCAAGTGA TGCTCCTGCC TCAGCCAACC AAAGTGCTGA
11701 GATTGCAGAT GTGAGCCACC ATTCCCTGCA GGAACAGTCT TAGATTTATC
11751 CACGTAGTCA CTGTTTCTGG TGCTCTTAAT TCCTTTGTAC AAATCCAGAT
11801 TTCCATCTGG TATAATTTTC CTTCTACCTG AAGGATGTTA TTTTCTTCTC
11851 TGTGTCAGGC CTGTTGGTGA TTAACCTTTT CAGCATTTT TTTTCTTTT
11901 TATTTGACAG GATTCACTCT GTCACCTCAG CTGGAATGCA GTGGTACAAT
11951 TATAGTTCAC TGCAGCCTTG AACTCCTAGA CCAAGCAAT CCTCAGCCTC
12001 CTGAGTGGCT GGGACTATAG GCATGCATCA CCACGCCTGG ATAATTTTTT
12051 ATTTTTACTT TTTGTAGAGA TGGTGTCTTC CTACGTTGCC CAGGCTGGTC
12101 TCTTACTCCT GGGCTCAAGG GATCCTCCA CGAGGGATCC TCCCACCTTG
12151 GCTTCCCAA TGTGTTGGAT ACAGGTGTGA GCCACTACTC CTGGTCTTCT
12201 TTCAACTTTT GACTTTTGTA TGTCTGAAAG TCTATTTTGC CTTCATTTTT
12251 CAAAGATAGT TTTGCTGGTT ATAGAATTCT AGACTTTTTT TTTCTTTTCA
12301 GTACTTTTTA AAAAAAAGTT ATTTATTATT TTTTATATT TGAGACAGTG
12351 TCTCGCTGTG TCACCCAGGC TGGAGTGCAG TGGCACGATC TTGGCTCACT
12401 GCAATCTCTG CCTCCTGGGT TCAAGTGATT CTCTGTCTC AGCCTCCCAG
12451 GTAGATGGAA TTACAGGCGC ATGCTACCAC GCCCGGCTAA TTTTGTACT
12501 TTTAGCAGAG ATGGGGTTTT GCCATGTTGG CCAGGCTGCT TTCGAATCC
12551 CTGACCTCAG GGTGATCCGC CCTCCTTGGC CTCCCAAAGT GCTAGGATTA
12601 CAGGCGTGAG CCACTTTGCC TGGCCTCAGG ACTTTTAAAG ATGCTATTCT
12651 GCTGTCTTCT GGCCCACTTC GTTCAAGAA GTGTGCTGGC ATTCTTTGGT
12701 CCTCTGTATG TAACCTTTTT ATCCACCGCT TTTAAATAT TCTTTATCAC
12751 TGGTTTTAGG CAGTTTCATA ATGTGGCTTG ATGTAGTGT TTTTGTGTTG
12801 TTTGTTTTTT GTTTCTGTG TTTATGCTTG TGATTCATTG AACTTCTTAG
12851 ACATGTGGGT TTGCAGTTTC ATCAAATTG GAAAAATAAC ACCCATCTTT
12901 TCGTGAAATC TTTGTACTGT CCCCTATGC GATTCCACAG ACTTCTTTG
12951 CAGAATCCCC TGTCTGTATT TCAGGCACCT CGAAGGTGTT CTCACAGCTC
13001 ACTTGGGATT TTCAGCCTTC TCCTTTGTGT GTTTCATTTT GGATAGTTTC
13051 TGCTGCTACA TTTTAGTCTT TTTTCTCTGT AATGTCTCAG CTGTTGTAAT
13101 TGCACCAAGT GCTTGTTCCT TCTCAGACAC TGTGGTTTTT ATTTCTAGAA
13151 ATCCAGTTTG AATCTTTTTT TTTCTTTTTT TTTTTTTTTA AGATGGAGTT

```

13201 TCGCTCTGTT GTCCAGGCTG GAGTGCAATG GCGCGGTCTT GGCTCACTGT  
13251 AACCTCTGCC TCCCGGCTTC AAGTGGT TTT CCTGCCTCAG CCTCCCAGGT  
13301 AGCTGGGATT ATAGGCATCC ACCACCATGC CCAGCTAATT TTTGTATTTT  
13351 TAGTAAAGAC AGAGCTTCAC TGTGTAGGCT GGGCCAGTCT CGAATTCCTG  
13401 ACTTCAGGTG ATTCAGCTGC CTCGGCCTCC CAAAGTGCTG GGATTACAGG  
13451 CGTGAGCTAC CGTGCCCGAGC CTCGAATTTGA ATCTTTTAAA ATGTATTTTC  
13501 TACATCTTTA CTTAACAGTT GCAATCTTTC CTCTGTTTTT TTTTAAAGTA  
13551 TATGGAATTT GGTACAGTT ATTCTTTTAA TGTCTTATTA ACTAAATTAC  
13601 ATTTGGTTAC AGTTACTCTT TTAATGTCCT GTTAATTCTG TCATCTTTAA  
13651 CTATGTTACT TTTGGGCTCT TTTTATTTT TTTATTTTGA TTTTGTGAGA  
13701 CAGGGTCTCA CTGTGTCACC CAGGCTAGAG TACAGTGGCA TGATCATGGT  
13751 TCACATATAGC CTCAACCTCC CAGGCTCAAG TTATCATCTC ACCTCAGCCC  
13801 CCCAAGTAGC TGGGACTATA GGCATATGCC ACCATGCTCA ACTAATTTTT  
13851 TATTTTTTGT GGAGACAGTG TCTCACTAAT TTGCTCAGGC TGGCCTCAAA  
13901 CTTCTGGGCT CAAGCAGTCC TCCTGTCTTG GCTTCCCAA GTGTGGGAT  
13951 TCAGGCTATG GGCCACTGCA CCCAGCCATA TTTTCTGCT TCTTTCGATG  
14001 CCTAGTAATT TTTGTTTGA TGGATGCTAG ACTTTGTGAG TTTGAGGTTA  
14051 TTGCGTGCTG GATATTTTTG TATTCTATAA ATCTTCTTGA GCTTTGTTCT  
14101 GGGATACAGT TAAGTTACTT AGGACAGTTC GATCCATTTA GGTCTTACTT  
14151 TTAAGCTTTG TGAGCTGGAC CAGAGCAGTG TTCATTCTAG GGCAAGCCTA  
14201 ATTTGGCCCA CCACTTGATT TTGTAATAA AGTTTTATTG GAAAATCACT  
14251 CCCATTCAAT TACATGGTCA ATGTCAGGTT TTATGCAATA ATAACAGACT  
14301 TGAGTAGTTG CGCCAGAGAC TTTATGGCCC ACAAAGCAAA AAGTTTTTAG  
14351 TATTTAGCCC TTTACAGAAA ACATTTGCCA ACCCCAGGCC TTGGGCTAAT  
14401 CTTCCCACT GCTGAGGTAA AACCTCTGG AGTCCCTGAT GCTCCATGAA  
14451 TTTATGAGGGT TTTTACTCT GGTGTGAATC CTGGGATTG TCCCTTGTA  
14501 CCCTCTTTGT GACTCTTCCC CACTGTGGGT TTGGGTCATT TTCGCCAGC  
14551 TTTGCCTCGA CCAGTTCTCT GCTGCACATC CGAGGGCACC TCTGCAGATC  
14601 TCTGCAGCTC TGTCTCCGGC TGCCCTCTCT GCTCCTGTGA CCTTGCGGTC  
14651 AGCTCCAGCT GCAGGCCTCC TCAGACTTCC TCCCAGCATT GCCCCCTGAA  
14701 CTCAGGGAAA CCCCTGGGCT CTGCTAGGGG GTCCCTCCA TACACATGGG  
14751 GACCATCGGG GGAACCGTA GGCAGCCGGC GGGGCACTGA TAGACTCACC  
14801 TCCCTTGTTT CCCTGTGGCC GTCATGGCTG CTGTCCCGCG TCTTAGTGGC  
14851 CATCATGTCC ACTGTCCCGC ATCTTGAGAG CCATTGTTT TGTGTTTGG  
14901 GAAAGGGGGT AAATCTGGTT TCTTCTCTCT TTGTGCTTTG AGGTGGATGT  
14951 GCCTGGGGCC ACCTGATTTC AGAGAGTCTT TGCCGCGGTG CACGATGTCC  
15001 AGTCACCTGG AGCTTGGCAG CAGGTGGCAT GTGCACCTGT CCGCAGCCTG  
15051 CAGCTCTGCC GTCCACCTG CTTATGCTGC CACATAGCAT TTATGTCTGT  
15101 GTTGTGTTTC CCAGAGTGCA GAGTGAGGAC TGTGGCCCGG TGGGGGCGTC  
15151 TCCTGCCTCT GGGATGCTG CCCCAGTAAC CTGGAGGCAG CCGCGCTACC  
15201 CCACACTCGC CGGGAGCAGG GTCTCGGACG CACCTCTCT TCTCTCTAG  
15251 CTCGCAACGC CCTGGTGGTC TCCTTCGCAG CCCTGGTTGC GTACTCCTTC  
15301 GAGGTGACTG GATACCAGCC TTTTCATCCTA ACAGGGGAGA CAGCTGAGGG  
15351 GCTCCCTCCA GTCCGGATCC CGCCCTTCTC AGTGACCACA GCCAACGGGA  
15401 CGATCTCCTT CACCGAGATG GTGCAGGTGG GCGGAGCCGG GAGGCAGGAT  
15451 GCGGTGGCTG AGGCTGCAGT GGCCCTGGC CTGGCTCCTA CCCTGATGTA  
15501 TCTGCTGGGT GCCAGGGGGT CTGAGGTCAG TTAGGACAGC TGAGTCTCA  
15551 GGAACGGACA TCTCAGTTAT TAAAGAATCC CAGGTGGAT GCAAACTCAG  
15601 CGAGCTCAGG GATGTCACGT TTGTGTTTCT GGGCGCTTCT CCTGTTTTGG  
15651 ACTCCAGCTG AGGATGAATT TACCGTGTTC CTCCAGCAC CTGGCGCCTC  
15701 TTCAGACAAG GAGGCGGATC CTGCAGCTGA CAAGCACTTG CTCCTGTTAC  
15751 CTGTGGGGCG GGGTGGGTCC TTGCTGCTTT CATGGGTCAC TGCTGGGTCC  
15801 TACCCCTTAG GAAGGTCACT CACCATCCCT CTCTCCTCTC TCAGGACATG  
15851 GGAGCCGGGC TGGCCGTGGT GCCCTGATG GGCCTCCTGG AGAGCATTGC  
15901 GGTGGCCAAA GCCTTCGGTA AGACGCTGT CACCCACACC CCAGGTCTCC  
15951 CAGTGCGCCG GCTGGGCTAG GCCTGCCTGC TTTCTAGCTT GCCTTTATCC  
16001 GTTACTAGTT TTAGAAATTT GAATTCATAT CCAAGTAATA CATGCTCATG  
16051 ATAGATACAT ATGTATTGTG TATATATGAT AAACTGGAT CTATAATGAG  
16101 GCATGCCCTC CCACCCCATG GTGTGCTGGT GAGTGTGTA ACAGCCTCTG  
16151 CTGTTTGTGG AAATAAAGG TTTTGCTTCG TGGCCCTTGC CGATGTCCAC  
16201 GGTGTAAACG CTGCTGTCTG ATTTTAAGGT AACGTCACTG AAAGGGGAGT  
16251 TTGCACATGG AGCTGGGTTG AGATCTGCAT GAACAATCAT ATTCTATGGT  
16301 GTCTCCACCA TGATAGATACA GTGGGTGCAA ATAACCTCAT CAGTAGTAGC  
16351 CAAATGCCAA ATAAATTAGG AAGTGATGAG TTTTAAAGTAT TATCTTTGGG  
16401 CCAGGCATGG TGGCCAGGC CTGTAATCCC AACACTTTGG GAGGCTGAGA  
16451 CGGGAGGATC GCTTGAGCTC AGGAGTTTGA AACCCACCTA GGCAACGTAA

16501 CGAGACCTCG TTTCTATTAA AAATAAAAAAT TAGCTGGGCA TGGAGCACAC  
16551 CTGTGGTACC AGCTTCTCAG AAGGCTGAGG CAGGAGGATC ACTTGAGCCC  
16601 GGGAGGTCGA GGTGGCATTG AGCTGTGATA GTGCCACTGC ACTCCAGCCT  
16651 GGGCAGCAGA ACAAGATCCT GTATTATCTG TTTAATTGAA AGTTTAAAT  
16701 TAATTGGTAA TAATGGCTAT GTTAGTAAAT AGGCTCACAA AACTCCTAAA  
16751 CATTCAGCAA CATGCTTTAT TCAGCTGGCT CAGCCCATCG GCCAGCCCCCT  
16801 CCTCGCCCCC AGGGAGGCAG GGAACCCCTC GTCTCCTTCA GTGTCTGTTT  
16851 TGAGTCAGCA TCTCTAGATT CCCTCCTTGC AGCTCCGTCC TTCACTCGCT  
16901 CCACTCCTGC CAGCTTTGTA TTGTCTGTGC GGATGACACC TTCGTTCTGT  
16951 CTTGTACCCG TCATTTAGTC CTCTTGCGT TTGGGTTTAT GTTGATTCTA  
17001 AAGGTTACAG CCCAGTTACC AGTGTATGTT ACTGTGACTG TGTAATACGT  
17051 GTTTACTGCC TAGCCCAGGC TGTGCCAGG TAAAGTTCC TCTCCAAAGT  
17101 CCCAGTGCTC TAACCCTTAC CCCCTTCAAA GGAGTTTTTT TTGTTTTTTT  
17151 TTTTTTTTTT TTTTGGAGA CAGAGTCTCA CTCTGTGGCC CAGGCTGGAG  
17201 TGCAGTGGCA CGATCTCTGC TCACTGCAAC CTCGCTTCC CAGGTTAAG  
17251 CAATTCTCAT GCCTCAGTCT CCCAGGTAGC TGGGAATACA GGCATGCGCC  
17301 ACCACGCCCA ACTAATATTC TATACTTTTA GTAGAGACAG GGTCTCACTA  
17351 TGTGGGCCAG GCTGGTCTCA AGTGATCTGC CCACCTGGGC CTCCCAGTAA  
17401 TTTTTTTTTT TTTTTTTTTT GAGATAGTGT TTCACTCTTG TTGCCCATGC  
17451 TGGGTGCAAT GGTGTGATCT TGGCTCACCG CAACCTCCGC CCCCAGGTT  
17501 CCAGCAATTA TCCTGCCTCA GCCTCCCAAG TAGCTGGGAT TACGGGTGTG  
17551 CACCAACATG CCTGGCTAAT TTTGTATTTT TAGTAGAGAT GGGGTTTCTC  
17601 CATGTTGATC AGACTGGTCT CAAACTCCTG ACCTCAGGTA ATCCGCCAC  
17651 CTCAGCCTCC CAAAGTGCTG GGATTACAGG TGTGAGCCAC CATGCTCAGC  
17701 CCTCCAGTAA TTTCTTTTTT TTTTTTTTTT GAGACGGAGT CTCGCTCTGT  
17751 CGCCCAGGCT GGAGTGCACT GGCGTGATCT CGGCTCACTG CAAGCTCCGC  
17801 CTCTGGGTT CACGCCATTC TCCTGCCTCA ACCTCCCGAG TAGCTGGGAC  
17851 TACAGGCGTC CGCCACCACG CCCAGCTAAT TTTTGGATT TTTAGTAAAA  
17901 ACGGGGGGTT CACCGGGGGT CTCGATCTCC TGACCTCGGG ATCTGCCCTG  
17951 CCTTAACCTC TCAAAGGGCT GGGATTACAG GCGGGAGCCC CTGGGCCCCG  
18001 GCCGTAATTT TTTAATGGAA AAACAAGGGC TCACCTTGGG GGCCAAGGCT  
18051 GACCTCAAAC TCCTGAGTTC AAGGGATCCT CCTGCCTCGG GTTTNNNNNN  
18101 NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNTGGTCT  
18151 GGGCTAATTT TTGTTTTTTT GTAGAGAGAG TATTTGCCAT GTTGCCAGG  
18201 CTGGTCTCGA ACTCCTGGGC TCAAGCGATC CTCCTGCCTT GGCTCTAAA  
18251 AGTGCTGCCA TTATAGCTGT GAGCCACCAT ACCTGGCCTG ATTACATGGT  
18301 ACTGTACCTC TTTCCACTCT GTGATACATT CAAGCACATG TTTACACGCA  
18351 CATATATGTG AATATTTTAG GTGTGTATTC TTAGATTAC AAAACTTATT  
18401 GCCGTGTGTA GAAATTCATA ATCTGTGTTA CATTTTGTGA ATACTTTTGT  
18451 AGCGATGTTG AATATTACTG ACCAGGTCTT TACCGTTGGT TCCCTGTAGC  
18501 ATCTCAGAAT AATTACCGCA TCGATGCCAA CCAGGAGCTG CTGGCCATCG  
18551 GTAAGACCCC AGCCCGGGGA AGGAAGACAC CAGCTGTGGG CCTCCAGGGT  
18601 CCCAGGCCTG CCCCTCTGTG TCTCCTGCAT TGTAGGAAGA CCATGATGGT  
18651 GGTGATGAAC TGGGAGGGCA AAGGTGGCCC CAGATGGGAT CTCTGGAAT  
18701 ATTTAGTTTT GTTTTGGGTT TTTGAGATGG AGTCTCACTC TGTGCCCAG  
18751 GCAGGAGTGC AGTGGCACGA TCTCGGCTCA CTACAACCTC TGCCTCCTGG  
18801 GTTCAAGTGA TTCTCCTGCC TCAGCCTCCC AAGTAGCTGG GATTAAAGGC  
18851 ATATGCCACC ATGCATGGCT AATTTTGTGA TTTTGAGTAG AGATGGGGTT  
18901 TCACCATGTT GGCCAGGCTG GTCTCGAACT CCTGACCTCA AGTGATCCGC  
18951 CCGCCTCAGC CTCCCAAAGT GCTGAGATTA TAGGCGTGAG CCACTGCGCC  
19001 TGGCCTATGC CTTGTATCT TAAACCTTGA GACTCAGAGT GTGGCCTGGG  
19051 ACCAGCAGCC TGGGCACTGG GCATCTCCTG GGAGCTTACT GGAAAGGCA  
19101 GATCTCAGGC CCCACCCAG ATCTCCGAAT CAGGATCTGT ATTCTTCAGG  
19151 GCACGCCCCAG GGGATTCATG GGGTCAGCTT AGAAGTCAA GTTTGGAAGC  
19201 CGGGCATGGT GGCTCACGCC TGTAATCCCA GCACTTTGGG AGGCCGAGGG  
19251 AGGCAGATCA CCTGAGGTCA GGAGTTTGAA ACCAGCTTGG CCAAACCCCG  
19301 TCTCTACTAA AAATACAAAA ATTAGCTGGG TGTGGTGGTG TGTGCCTGTG  
19351 ATCCAGCTA CTTGGGAGGT TGAGGCAGGA GAATCACTTG AACCCGGGTA  
19401 GCGGAGGTT ATAGTGAGCC AAGATTGCAC CACTGCACTC CAGCCTGGCA  
19451 ACAGAGCGAG ACTCCTTCTC AAAAAAAGT AAAAAAAGT CAAAGTTTGG  
19501 GAAATGCTTA GAGACCCCAT GTTTTTCAA GACTTGTTTT GTATGACTTT  
19551 ATGAGATGAT CAAGCAGGTT TGGCCACAAG GGGAGACAGC CAAAGGCTCA  
19601 GGAAGATAAC CAGTTTTTTT TACTCAAAG GTCCCTTAGT TACCAGGGAG  
19651 GCCACAGCAT GCCTACTTAG GCCCATGTCA GAGGACACCA GGGGTGTCCC  
19701 CGGAAGGATA AATCCCAATC TGTGATTTA GGCTCCGCC CTTTTTCGGG  
19751 AATTTCTTTT GCTTAATGCT AATTTCAATA AAGGCCCCCG GCTTAAGACT

19801 TTTCATATGT TATTAATTTT AGGCGGGTTC TAACTCCTTA ATGAAGAAAT  
19851 ATTCTCCCTT CCCTGGATTT CCGAATCACA CCATTTCTTC GAGAAACGCC  
19901 TCCCACNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN  
19951 NNNNNNGTTT TTTTTTTGGG CACCCGGCCA TCATCTGGGG AGGCAGCATT  
20001 AAGGCCCATC ACTTTGCCCT TCAATGTGAC TGTCTCTTTT TTACACGAAA  
20051 AAGTTTTTTG GTGGAACCC GGTGGTCGGC CCCATTACCT CCTGAGTGCC  
20101 TAGGGTTTGT GTTCCGGGCC ATTTATCTCA GCTTTCACCC CCTCCTGGAT  
20151 GGTAAAGATG TCCCTCTGT TCTGCGTGGG GAAGGCGGGT CCTGGTCCCA  
20201 CAGGATAGAT TGGGTCAAAA GTGTTTCAGA GCTAGGATCT CTTCAGGCT  
20251 CTCGTGGGTC ACTCCTGTTT CCCAGCACTT TGGGAGGGTA AGCCGAATGG  
20301 GTCACCTAAG GTCAGGCATT TGAGACCAGA TTGGCCAGCA TGGCACACTC  
20351 CCTGTGTGCT ACTAAAAATA GAAAAATTAC ACTGGGTGTG GTGGCACATG  
20401 CCCGTAGTCC CAGCACTTTG GCAGGTTCAG CCGGGTGGAT CACTGAGGTC  
20451 AGGAGTTTGA GACCAGCCTG GCCAACATGG CAAACTCTG TCTGTACTAA  
20501 AAATACAGAA ATTAGCCAGG CGTGGTAGTG TGTGCTGTGA ATCCCAGCTA  
20551 CTCAGGAGGC TGAGGCAGGA GAATCGCTTG AACTGGGAGG CGGAGGCTGC  
20601 AGTGAGCTGA GATCGGCCCA CTGCATTCCA GCCTGGGCAA CAAAGCCAGA  
20651 CTCTGTCTCA AAAAAAAAAA AAAAAAAAAA AAAGGCTAGG CTTTACGTCT  
20701 GCAAGAATGT GGCCTGTTTT CTTCCTTCCC TGAAGGAGTG CGTAGGCCCA  
20751 GGCCCCCAGG ATGATTCTCC CGAGCCCTGC CTTCCTGCCT CCCTTCTGGG  
20801 TTTTGTGCTC CCTCCCATC TCCTTCCCT CCATCCTGTG TGCCTTCCCT  
20851 CCACGATCAG CCTGTCTTGC CTCCTCCCA GGTCTACCA ACATGTGGG  
20901 CTCCCTCGTC TCCTCCTACC CGGTCACAGG CAGCTTTGGA CGGTGAGTGA  
20951 CATGTCCGCC TCTTCTGTTT GCCCACGTTG GACGCTTAA CGTTGTACG  
21001 CTGACAAGGA GTCTGCCTGC CCTGACCCCG GCGCCCCGTC CTCCACTGTG  
21051 AACGCTCCGT GGAGAGGCAG GGCTGGGGGT CACCCACTGT CCTCCAGGGT  
21101 GTTCTCTGTT TCTTTATTCT CATAGATCGT CCTGCAGTTT CATACTAGAA  
21151 AGTTCCACTG GGCATTGTGG TACACCCTTG TTATCCCAT TACTCTGGAG  
21201 GCGGAAGCAG GAGAATCGCT TGAGCCAGC AGGATGAGGC TGCAGTGAAT  
21251 TGTGACCGCA CCGCTGCACT CCAGCCTGGG CAACAGAGCG AGACCCTGTC  
21301 TCTAAAAATA ATACTAGAAA GTTCCAGCA CGCCAAAGCC CTCCTAGCTC  
21351 CTGGTGCCAG AGTCAATTCC TGAAGGACG TGGAGATAGG AAGGGCCTCG  
21401 GCTCTGTCTT GAAGCAGCCG GGCATGAAGC TTAGCCAGA TGCCCTACGG  
21451 CCCCTCTCA GTCCAGACAA CAGGATGGAG GTGACCTGTG GCTTAAAGGA  
21501 GAGAAGGAGG CGTCGCCTGG CACTGCCCAG TCCCCAGCT GGTGACCCTT  
21551 GCCCTGCTGG GTATGGGGGC CCCACCTGGA TGGGGGAGG AGACAGAGTC  
21601 GGCAGGAACC TGAAAGGACA CGTGCTTCCT GAGCTTCTTC CTATAGTCAG  
21651 GGTGGCCCAA GCGCGGCTGT CTGTGACTGC ACCCTAAGTC TCTTTGCCTC  
21701 GGTCCCCCTG CAGTCCCCGC CTGCTTCCA AGCCGTGCTG GGAGCTGACG  
21751 TCCCCTCGGA AGATCAGCCA CAGGAGTGTG GACTGAGGTC TCCCTTTTCC  
21801 CGGCCCTG TGACTGACGG TCTCTGTGTT GCCTTCCAG ACAGCCGTGA  
21851 ACGCTCAGTC GGGGGTGTGC ACCCGGGCGG GGGGCTGGT GACGGGTAAG  
21901 GCCCCCATC TTCCCCTGT GCGCGCAGCC CTGAGAGTGG GAGAAAGGGA  
21951 GGAGGGGGCC CACAGAGACG TCCCTTTGGC TCATGGGCGG TGCGCCCCGG  
22001 GACTGCACAG GGAATTGGGG GGCCACACAG GAGTAGGGGG ACCACAGGAG  
22051 ACTGAGCAGG GGCTGGGGGC CTTGGCAGTC GTCGCCCTAC CCCCACCCCT  
22101 GTCCCCAGTG GGCTCTGCTG AACAAGAGGC TGCTACGCTG CGTGCTGGGG  
22151 GGACCTGCA CTCCCAGGT CACCTGTGTT CCGTGCCCC GCAGGAGTGC  
22201 TGGTGTGCT GTCTCTGGAC TACCTGACCT CACTGTCTTA CTACATCCCC  
22251 AAGTCTGCCC TGGCTGCCGT CATCATCATG GCCGTGGCCC CGCTGTTTGA  
22301 CACCAAGATC TTCAGGACGC TCTGGCGTGT TAAGAGTACG TCCTTGTCTT  
22351 ACAGGGGAGA GCGCTGTGAT GCGGTGTCTG AACGCGGAGG GTGTCATTTA  
22401 TGCTACCCCA TTTTCTGCA GCCCCTCTG TGGGGCTGGG ACTGGGAAGT  
22451 TAGGGCAGTC CCGGAACAGA GAAGTGATG GCCAGGAGAT GGCCCCAGAG  
22501 ATGGTCCCGA GGCTCAGTGG GAAGAGCTGG AGCTCCTTGT CTGACACCT  
22551 GGGGTCTTGA GCGGAGCACT GACCCGGGGG AGGGTCCCC CTGATCCCC  
22601 CTGCCCCCAT CCTACCTCT CTTGCCACCC GCCTCCAGCC ACCACTCTGC  
22651 CCGGCCCCAG TGGGGGGAGG GACAGGAGAC GTCCCTGGTG ACCAGCAGGG  
22701 CCAGCGGAAC AGCCTTGCAC CCTGGCTCAG AATGGCAGTT CTTTTTTTTT  
22751 TTATTATTAT TATTTTATT TTTTTTATT GATCATTTCT GGGTGTCTTCT  
22801 TGCAGAGGGG GATTTGGCAG GGTCTAGGA CAATATGGGG TTGGGGGTAA  
22851 GGTACAGAT AACAGGATCC CAAGGCAGAG GAATTTTCT TAGTGAGAA  
22901 CAAAATGAAA AGTCTCCCAT GTCTACTTCT TTCTACACAG ACACAGCAAC  
22951 CATCCGATTT CTCAATCTTT TCCCCACCTT TCCCCCTTT CTATTCCACA  
23001 AAACCGCCAT TGTCTCATG GCGCGTCTC AGTGAGCTGT TGGGTACACC  
23051 TCCAGACGG GGTGGTGGCC GGGCAGAGG GCTCCTCACT TCCAGTAGG

23101 GGC GGCCGGG CAGAGGCGCC CCTCACCTCC CGGACGGGAG CAGTTCCTTT  
23151 AACTTAACAC ATTTTGT TTTT GTTGTGGAA GAAGCATGGA GTGGGTTTAC  
23201 CTAGCTTACT TGGCATT TTTT TGGTTAGCCC CAGCAAGTTG CCAGGTGAAA  
23251 GGATGGATAA TTTTCTTGCA TGCCCGTCGC ATGCCAGGGC CTTTCGCGTG  
23301 CCAGTGCCTG GGGGCTTTAG AACAGCCCTG AAACGTGTTC ATGTGCCGGC  
23351 GTGGAATGGG ATGGCCGCTG CTGCTAGAAA CCAAGGCTCA GCCAGGGTGG  
23401 GGCAGGCCTG ACCTATGCGT GCGGTGGAAT CCCCCACAGG GCTAAGCCCC  
23451 TGCACTTTGT CCCCAGGGCA CCTTCTCCTT GGCCAGGTCT CAAGGGCTCA  
23501 CGTGGTCCCT GCCCCACTCC TCAGGCCAGC TCTGTGCCCT GACAAGCCCC  
23551 TGCTGCTGCC CTCCTGAGG TTGGAGGCCA GGGACCGGCC GGCAGGTCTC  
23601 ACCTGCGCTC AGCTCAGATG GGGAGGGCAT TTCTTTCTTT CGACTTGAAG  
23651 CATGGCCTGG TCAGCAGCTG CTGTCCCAA GTCCCTCAGG GCTGCTTGGG  
23701 GTC CATGAGC ACCTTTACTC ATATGTGGGG GGCAGAAGGC TGTCCCGCTG  
23751 TGCAGCAGGG CCATGTTGGG GCCTCGGGCA GCTGCCGGGC ATTCTCAGC  
23801 TGTGCCCTTC TCCTAGGGCT GGACCTGCTG CCCCTGTGCG TGACCTTCCT  
23851 GCTGTGCTTC TGGGAGGTGC AGTACGGCAT CCTGGCCGGG GCCCTGGTGT  
23901 CTCTGCTCAT GCTCCTGCAC TCTGCAGCCA GGCCTGAGAC CAAGGTACCC  
23951 CTCCGTGGCC TCTGAGTGGG GAGTGTGCTG GGGGCAGGAT TCCTGGGCAT  
24001 GGTCTTATGT TTTGAGGGTC CGGGGTGATT GTGGTCTGGG GTGCTGCTGA  
24051 AGGGGACCGC TCGTGGCAG GTGGGCAGTC ACCTTGCTAT AAACCATGGT  
24101 GTTCTCCAC TGTGTGGGG CCCTGGGGGT CTCCCTTAG CACCCCTCTC  
24151 CCGGTCCCTT GCAGCAGCT AGGTGGGGT GGGGCTTCCC GCTTGGGACA  
24201 GGCCAAGCCT GGTGGAGGCC ACCCGGTGAG ACCCGCTCC AGGACTCACT  
24251 CCTCCCCACA GGTGTCAGAG GGGCCGGTTC TGGTCTGCA GCCGGCCAGC  
24301 GGCCTGTCTT TCCCTGCCAT GGAGGCTCTG CGGGAGGAGA TCCTAAGCCG  
24351 GGCCCTGGAA GGTGCATGGG CGGGGGTCAA GGTGGTCTGA GGTCACTCCC  
24401 CTGTCTCTG CCCCCACTC CCTGCTGTTT AGGACCCCAA GACCCTGTCC  
24451 CCGACGCTCT CCAGTCCACA AGGATGCAGG CATCTCTGAG TGGGCTGGAC  
24501 CGTCTCTGT GGGCTCAGC CAGTGGCTGC TGCAGCAAGG GTGGTGGCTC  
24551 CCCACATATC ACTCCTTCCC TGCCCTAAA GTCCGGTTCC TGTTTCTGGG  
24601 GGGTTGATTT TAGGGGAGCT AAGGGCCTGT GAGTCTTAGG AGGGAACAG  
24651 CTGCTGCTGT CACCAAAACA TTGTCTCTGG TCCTGCCACC CGAATCCCCC  
24701 AACTGGGCGA CTCAGCCGCC ACGAGATGGA GCACTCTGGC CTCTCTGTCC  
24751 CCTGCCCCTG GCCAGAGCCT CCTTGGCCT CTGCAGAGCA GCTTTGGGCT  
24801 GCTCTGGGTG GCGTGACCTG GCTCGGGCCT GTCTCCCCAG TGTCCCCGCC  
24851 ACGCTGCCCTG GTCTGGAGT GCACCATGT CTGCAGCATC GACTACACTG  
24901 TGGTCTGGG ACTCGGCGAG CTCCTCCAGG ACTTCCAGAA GCAGGGCGTC  
24951 GCCCTGGCCT TTGTGGGCCT GCAGGTGGGT GTGCACTGGG ATGCCTTAGG  
25001 GGT TAGCAGC TGCCGGAAGG CCTTCTGTG CCTGCCCTCC ATGGCGAATG  
25051 TGACATCTCT GGGCTGTGAT GCTGGACGGC CCTTCGGCCG GTGCTGGCTC  
25101 TGCTTCTGAT TTAAACAGTT CTTGTCCCA TCTGGCCTTC CTCGTCCCTC  
25151 CCTGTGGAAG GGGGAGCGGT GGCCCCCAGC CCTCCGAGG GTCACGTTAT  
25201 GGCTTCTGGT CACTGCCACA CTGTCTTTG TGCTGGGGAC ACACAGTGAA  
25251 CGAGGGTCAG TCCCTGCTCT CAGGGGAATT GTATTTTAGG AAACAAAAG  
25301 AGACAGGTGA GGCTAGGCGT GGTGGCACAC TCCTGGAATC CCAGCACTTT  
25351 GGGAGGCCAA GGC GGCGCAGA TCACCTGAGG TCAGGAGTTC AAGACCAGCC  
25401 TGGCCAACAT GGTGAAACCC CGTCTCTACT AAAAATACAA AAAAACTAG  
25451 TCAGGTGTGG TGGCGAGCGC CTGTAATTCC AGCTACATGG GAGGCTGGGA  
25501 GAATCGCTTG AACCTGGGAG GTGGAGGTTG CAGTAAGCCG AGATCCCACT  
25551 ACTGTACTCC AGCCTGGGCG ACAGAGTGGG ACTCCATCTC AAAAAACAAC  
25601 AAACGAAAAA AAACAGGTGC TCATAGAATT TCATGAAAA CGTATTGTCA  
25651 GGGCTTCCAG AGGCTGAAGA CGGGTTTCTA TGGAGGCCGT CCTGTTTCTA  
25701 GCCGAGGTA AAGTGAAGG GCTGGGTCCC AGGCCCTGCG TCTTAGGCCT  
25751 CACCTAGGAG CTTTCTGAGC ACTGCAGGGT CAACATCCCA GGGGTGTGGC  
25801 CAGTGTGTTG AGAGAGGCAG GGGTCTCTGT TGCTGTGGTT AAATGTGCGC  
25851 TCTCTGCCAA GTCCTAGATG GCAGAACGTG GGGGACTAGG GCGTGTCCCC  
25901 AGGCGCCCGAG AGGAGACAT CATGAAC TAGGAGGCAAG GAGAGCCAG  
25951 TGACCCGTAT ACCCCAGGTG TGGACCACAG CCGACCCCTG TCAGAGTTTC  
26001 CTTCATTCCC AACCTGGCCC ACGAGGCTAG TGTATCTTC ATCCCCATGT  
26051 CCAGAAGAGG TCACTGAGGC CCAGAGAAGC CAGGTAATCT GCCCACGGTC  
26101 ACACAGGATG GTGGGCCAG CTTAGATTT GGGCATGTGC CTTAGAACT  
26151 TCGTCTATAA GTGTTACGTC CTGTGTCATA TTTGAGAGC ACATGTCTTC  
26201 ATGGTCTTTG GAGATGATTC ACTTAAAAA ATACTCTCT GACCCAGACA  
26251 TGGTGGCTCA CACCTGTAAT CCCAGCACTT TGAGAGGCCG AGGCGGGAGG  
26301 TCAAGGGTTT GAGACCAGCC TGGCCAACAT AGTGAACCTC CGTCTCTACT  
26351 AAAAAACAAA AAATTAGACG GGCATGGTGG CGCACGCCCTG TAATCCCAGC

26401 TGCTCAGGGG GCTGAGGCAG GAGAATCGCT TGAACCCGGG AGGCAGAGGT  
 26451 TGCAGTGAGC CGAGATCGCG CCATTGCACT CCAGCCTGGG TAACAAGAGT  
 26501 GAAACTCCAT CTCAAAAACA AACAAACAAA CAAACAAAAA GTCCTCTGGG  
 26551 CCAGGCGCTG TGGCTCACAC CTGTAATCCC AGCACTTTGG GAGGCCGAGG  
 26601 TAGGAGGATC ATTTGAGGCC AGGAGTTTGA GACCAGCCTG GGCAACATGA  
 26651 CGAAACCCCA TCAAAAAAAT AGGAAAAAAT AGCTGGACGT GGTGGTGTGC  
 26701 ACCTGTAGCT ACTCAGGAGG CTGAGGTGGG AGGATCACTT GAGTCTGGGA  
 26751 GGTGAGGCT GCAGTGAGCC ATGATCGTGC CACTGCACTC CAGCCTGAGC  
 26801 AATGAGCAAG ACCCTGTCTC AAAAAACAAA AAATTTAAAA AAAAAAATC  
 26851 TTCTCTGAC AGCATTCCCC TGGGGCTGCG TTTCTTCTCA CCATTCACTG  
 26901 GTATGGAGGT GAAGCCATAC CTCTCCGGGA GACTCTGAGA TGGCATGTCT  
 26951 ACCAGGCTGC CGACCCGTGT GCTACAGAGG AACATCCCTG CCCTGGCTAA  
 27001 AGTCTGTCTG TCTCTCAGGT CCCCGTTCTC CGTGTCTGCTG TGTCCGCTGA  
 27051 CCTGAAGGGG TTCCAGTACT TCTCTACCCT GGAAGAAGCA GGTGGGCACA  
 27101 GTCAGACATC CTGTGGCTTT GGTGATTTTG TAAAAATCAT AAATGCTTAT  
 27151 TGTAATAAAT ATGGGAAACC AGCTGGGCAC AGTGCTCAT GCCTGTAATC  
 27201 CCAGCACTTT GGGAGGCCGA GGCAGGTGGA TCACCTGAGG TCAGGAGTTC  
 27251 GAGACCAGCC TGGCCAACAT GGTAAACCT CATCAGTACT AAAAATACAA  
 27301 AAATTAGCTG GGTGTGGTGG CATGCACCTC TAGTCCAGC TATTCGGGAG  
 27351 GCTGAAGCAG CAGAATCGCT TGAACCCAGG AGGCAGAGGT TGCACTGAGC  
 27401 CGAGATTGTG CCTCTGCACT TCCAGAGTAG GGTGAGAGAG CTAGCCATGT  
 27451 TGTCCAGAAA ANNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN  
 27501 NNNNNNNNNN NCCTGGTTGA GCAGGTATAA TCGCTTGCAA CCCAGGTAGC  
 27551 AGCAGTTGTA CTGAGTCCGA GAATTGTGCC ACTGCACTCC AGCCTGGGTG  
 27601 AGAGCAGCGA GACTCTGTC TCAAAAAAAA AAAAAACAAA AAAAAACAGG  
 27651 AACCACCTCC ACCCCAGTCC ACTTTGGTGA TCACTCCATA CCCCTCCCCT  
 27701 AAACACGCAT ATGTACCTGC CTGTCAAGAT GTGGATATAT GTTTGCGGTT  
 27751 TTACGTAAAT GGGACCAATT CATACCTGGT GCTCTGGAAC CCACATTTTT  
 27801 CATGCAGAAG GTTGGAAGGA TGTCTTCCA GCCGAAAGTC CACATCCCGG  
 27851 GGGATCAGGA CAGAGCAGGG CCGGGTCAGG AGATCCAGAA GCCCCGGGAC  
 27901 AGAAGGTACG GGAGGGACAG GAGCAGGGTG GGCCTGACC CTTGAGACAG  
 27951 CAATCGCAGG AGGTCTGAGC CGCAGCAGGT GTCAACAAGA GGATGGGCCA  
 28001 GAGATGCAGA GCATCCACCC CAGGCCACAC AGCAGTGGCC AGAGGGTCCC  
 28051 AGGCCCCAGT GCTAGGCCTC TTCCTCTTCC ACTGAGGTCA CAGCTGAAGC  
 28101 TGGGTCAGCT CCGTGAGAGT GAGGGGTGGC GGATGTTGTA CTGACTTCCT  
 28151 TGGCTCAATG TGACGTCAGG GAGATTCACC CATGTTGTTG TAGAATCAGC  
 28201 TCAGATGCAG TGCACTGGAG GTGGATAAGC AGAATGTGGC CTGGCCGTGG  
 28251 GATGGGAGGG TACCCTGGAG CAGTAAGAAA GGGCCGTTAG TCACCTGAAA  
 28301 AATACGCTTA CAGAGACTCA GGTGAGACCC TCATGGAGTT AGTGACACTG  
 28351 GCCTGGGTGG CCCACAGCTC CTTCTGTCAC ACCTTCCAGG ACTCTGGAAG  
 28401 GCCCTCCTTA ATCCCTTCTT GTGAAGTAC CCATCCTCAC TTCTGAGCTT  
 28451 TTAGTGCTTG AAACATTTAT TGTATTTTCT GCAGAGAAGC ACCTGAGGCA  
 28501 GGAGCCAGGG ACCCAGCCCT ACAACATCAG AGAAGACTCC ATTCTGGACC  
 28551 AAAAGGTTGC CCTGCTCAAG GCATAATGGG GCCACCCGTG GGCATCCACA  
 28601 GTTTCAGGGG TGTTCGGGAA GGTTCCTGTC ACTGTGATTG GATGCTGGAT  
 28651 GCGCCTGAT AGACATGCTG GCCTGGCTGA GAAACCCCTG AGCAGGTAAC  
 28701 CCAGGGAAGA GAAGGAAGCC AGGCCTGGAG GTCCACGGCA GTGGGGAGTG  
 28751 GGGCTCACTG GCTTCTGTG GGATGACTGG AAAATGACCT CGCTGCTGTT  
 28801 CCCTGGCATG ACCCTCTTTG GAAGAGTGGT TTGGAGAGAG CCTTCTAGAA  
 28851 TGACAGACTG TGCGAGGAAG CAGGGGCAGG GGTTCACAGC CCGGGCTGTG  
 28901 CGAGGCATCC TGGGGCTGGC AGCACCTTCC CGGCTACCA GTGCCACCTG  
 28951 CGGGGGAGGG ACGGGGCAGG CAGGAGTCTG GGAGGCGGGT CCGCTCCTCT  
 29001 TGCTGCGGC ATCTGTGCTC TCCGAGAGAA AACCAAGGTG TGTCAAATGA  
 29051 CGTCAAGTCT CTATTTAAAA ATAATTTTGT GTTTTCTAAA TGGAAAAAGT  
 29101 GATAGCTTTG GTGATTTTGT AAAAGTCATA AATGCTTATT GTAAAAATA  
 29151 CAGGAAACCA CCCCTCACCC TGTCCACTTG GGTGATCATT CCAGACCCCT  
 29201 CCCCAAACAT GCATATGTAC CTGTCCGTCA GTGTGTGGAT GTATGTTTAC  
 29251 AGTTCTACAT AAATGGGATC ATTTTATACA TGGTGTCTG GAACCCACAT  
 29301 TTTTCATGCA GTCATTTGCA GTGAATTATT TATTGTGATA ATAAATAGCA  
 29351 TTAGAATACA AGATTTTAA TGTCTGCGTG GTATTTTGGT CTATATATGC  
 29401 ATCATAATTG ACTTACCGAG CCCTCTGTTT AACGTGTGCG TGGGTTAGAG  
 29451 ACGGGATCGT GCCTCCTTTA GATGTGTCAG TTGAAGCCCT TGCCCTCTAT  
 29501 GTGACTGCGT TTGGAAACAG GGCTTTTAGG GGGTAACTAG GTTTAATGAA  
 29551 GGTCATAAAG GTGGGCCCTG AACCGATAGC TTTAGTGTCC TCGTGAGAAG  
 29601 AGACGCCAGA GAGCTCGTTC TCTGCACCT CACACCCAG GGAACCTCAT  
 29651 GAGAGGACAC GGCAAAACCA GGCCATGCGC CCACCAGGAA GAAAGGCCTC

```

29701 ATGAGGACCC CAGCCTCCAA AACTGGGAGA AGATGAATCT CTGTGCTAGG
29751 CCCCGCAGCC TGGGGTGATC TGTGACGGCG GCCTGAGCAG GTGAGGACTG
29801 CCTGCATGTT TGTATTTTAA TGAATGCTTT GATTGAGTCT GGGGGTAAAT
29851 CCCTGGAGGC CTGTGGCAGC CTCAGAGGTG TGTTCCTCCT GCACTTTCTT
29901 CAAGAAGAAT GTGGCTGCC CTGCTGAGCC TCGTTCGCCC CGTTCGCCC
29951 GGGCAGTCCC GGCCAATGTC AGCGCAGCAA GGGGAGGGCC TCTGTAACCA
30001 GGGCTGCTGG CTGCGGGGCT CCCCCTGGA CACGGGAGCG GACATTGGAG
30051 TGTCCCTTCAT CCGCGTCACT CTTCAGGTC CTTGCTCC CATTTTCCTT
30101 CCTTCCTTCC TCTTTCTTTC CTTCTTCCT TTCTCCCTT GCTCCCTTCC
30151 TTTTCTTTCT TCCTTCGCTT TTTCTTTCCT TTCTTCCTTC CTTCTTTTTT
30201 TTCCTTCCCG TTCTCCCTTT CTTCCTCCT CCACCCCTTT CTTCTCACTG
30251 TGTTCCTCCAG GCTGGTCTTC AACTCCTGGG CTCAAGCAAT CCTCTTGCCT
30301 CAGCCTCTGG TGTGGCTGGG ACCACAGACA CATGCCACCG CGCCAGGCTA
30351 ATTTGTTAAT TTTTTTTATA GAGACAGGGT CTCACCTTGT TTCCAGGCT
30401 CGTCTCAAAC TCCTGGGCTC AAGTGATACT CCAGCTTCAG CCTGAAGTGC
30451 TGGAATTAAG GTGTGAGCCA CCATGCCTGG CCCTCTCTCA TTCACAAGTG
30501 AACCGTTTAC CCCTGCCCTC CAATCCATGT CGTTTCTGAC CTCAGGCAGC
30551 TTCTCTTTCT ACATAAAGTG AACCTGTCCC AAAGCTGTGC TCACTGGGCC
30601 CCCCTGCCAG GGCTGGAAGA GGCAGCAGTT CACATTTGGC TTGCACTCAC
30651 ATACCAAGGG CATCCATGCT TGAACCTAGA CATGGTTCAT TCACAGGGAT
30701 GGGTGAAGGT AAACAGACTG TGGCGGGCAC TAGGCACTAT CAGTTTCATA
30751 AATCTGCATA CCACCTGTGA CTCAGCAGTT ATGCTTCTCG GAATCTACTC
30801 AAACATGCTT GAGTCAGCCT TCCAGGAAGC AGGCACAAGG ACGTTTGTGA
30851 CAACTTGGTC AGTAATTTTA AGAAGTAGGA AGAAACCTGA GTTTCTCTGA
30901 ATTCCGTAAC ATCTTGACTG TAGGACACAC GAATAATGCC GTGGAATATT
30951 GTGCAGCAGT TAAAAGAAAT GAGGTGAAGG CCGGGTGTGG CAGCTCACGC
31001 CTGTTATCTC AGCACTTTGG GAGGCCGAGG CTGGTGGATC ACTTAAGGTC
31051 AGGAGTTTGA GACCAGTCTG GCCAACATGG CGAAACCTG TCTCTACTAA
31101 AAATACAAAA ATTAGCTGAC CTCTGTGACA GAGTGAGACT CTGCCTCAAA
31151 AAAAAAAAAA AAAAAAAAAA AAAAATGAGG TGAAGCTTTA AGTTGTAACA
31201 CTGATTTTGG GGTGCAATAA AGCAAGTTGC AGAATGATAC CCATGTTAAG
31251 ATGCTATTGG AGTGAACACA CGGACCAAAC AATTCTATGT TGGGTACAAA
31301 TTTAAGAAAG AGTTCGGGC TGGGCACGGT GGCTCACAAC TGTAATCCCA
31351 GCACTCTGGG AGGGTGAGTC GGATGGATCA GGAGGTCAGG ACTTCTAGAC
31401 TAGCCTGGCC AATATGGTGA AACCCCATCT CTAATAAAAA TACAAAAATT
31451 AGCCAGGCGT GTTGGCGCGT GTCTGGAGTC CCAGCTACTC AGGAGACTAA
31501 GGTAGGAGAA TCACTTGAAC CCGGGAGGCA GAGGTTGCAG TGAGCTGAGA
31551 TCATGCCACT GCACTCTGGC CTGGGCAACA GAGCGAGACT CAAAAAATAA
31601 ATAAATAAAT AAATAAAATA AATAAGAGTT TTCTGCACTT TGGGAGGCCT
31651 GTAGTCCCAG CTACTCTGGA GGCTGAGGCA AGAGGATCAC TTGAGCCTGG
31701 GGGGGTCGAG GCTGCAGTGA GTCCTGATTG TGTCACTGAA ATCCAGCCTG
31751 GGCAACAGAG TGAGAC (SEQ ID NO:5)

```

## FEATURES:

```

Start:      914
Exon:       914-1147
Intron:     1148-2008
Exon:       2009-2286
Intron:     2287-2517
Exon:       2518-2563
Intron:     2564-5066
Exon:       5067-5146
Intron:     5147-7047
Exon:       7048-7190
Intron:     7191-15250
Exon:       15251-15426
Intron:     15427-15844
Exon:       15845-15917
Intron:     15918-18499
Exon:       18500-18550
Intron:     18551-20881
Exon:       20882-20942
Intron:     20943-21839
Exon:       21840-21895
Intron:     21896-22194
Exon:       22195-22335

```

Intron: 22336-23816  
 Exon: 23817-23944  
 Intron: 23945-24261  
 Exon: 24262-24361  
 Intron: 24362-24840  
 Exon: 24841-24974  
 Intron: 24975-27018  
 Exon: 27019-27091  
 Intron: 27092-28484  
 Exon: 28485-28576  
 Stop: 28574

## CHROMOSOME MAP POSITION:

#	SHGCNAME	CHROM#	LOD SCORE	DIST. (cRs)
1	SHGC-56719	17	15.9	14
2	SHGC-58932	17	15.68	14
3	SHGC-97	17	14.5	17
4	SHGC-53147	17	13.3	22
5	SHGC-33067	17	12.52	23

## ALLELIC VARIANTS (SNPs):

POSITION	Major	Minor	Context
30344	g	a	aagcaatcctcttgccctcagcctctggtgtggctgggaccacagacacat [g/a]ccaccgcgcaggttaatttgttaatttttttatagagacaggtctca
31170	a	g	ctctgtgacagagtgcactctgcctcaaaaaaaaaaaaaaaaaaaaaa [a/g]aaaatgaggtgaagctttaagttgtaacactgattttggggtgcaataaa
16256	c	t	aaacgctgctgtctgattttaaggtaacgtcactgaaagggagtttgca [c/t]atggagctgggttgagatctgcatgaacaatcatattctatggtgtctcc
13376	a	t	catgccagctaatttttgatttttagtaaaagacagagcttcactgtgt [a/t]ggctgggcccagctctcgaattcctgacttcagggtgattcagctgcctcggc
12210	t	c	atgttgggattacaggtgtgagccactactcctggtcttcttcaacttt [t/c]gacttttgatgtctgaaagtctattttgccttcattttcaagatagt
12072	g	c	catgcatcaccacgcctggataattttttactttttgtagagat [g/c]gtgtcttctacgttgcccaggctggtctcttactcctgggctcaagga
11922	g	t	taactctttcagcatttttttttttttttttttttttttttttttttttt [g/t]tactcaggctggaatgcagtggtacaattatagttcactgcagccttga
11903	-	a	ttgcaggcctgttggtgattaactctttcagcatttttttttttttttttt [-/a]tttgacaggattcactctgtcactcaggctggaatgcagtggtacaatta
10009	c	g	ctttgacctgtggattatttagaagtggtatgatgttcttctgtgaa [c/g]catgggagttatttagtttccaaatagttgggggtttcccaattagtact
4519	c	t	cctgccagagacctcagagctggtatcgtggtgggaggctcctactttgc [c/t]gaggattccccagctggtttcttgaagccctcagagccctccacatct
4181	a	g	aacagtcccagagtccgccccccaggatgtcctaaccaccaaccacctgt [a/g]cgcagcagttttagggcagggtccttgagcctgtggccatgggatccagg



20952	a	c	tccctcgtctcctcctacccggtcacaggcagctttggacggtgagtgac [a/c]tgccgcctcttctgtttgccacggttgacgccttaacggtgttacgct
20987	t	c	tggacggtgagtgacatgtccgcctcttctgtttgccacggttgacgcc [t/c]taacggtgttacgctgacaaggagtgctgcctgcctgaccccgccgcccc
21620	g	a	ccccacctggatgggggcaggagacagagtcggcaggaacctgaaaggac [g/a]cgtgcttctgagcttcttctatagtcaggggtggccaagcgcggtgt
21795	t	c	ctgacgtccctcggagatcagccacaggagtggtgactgaggtctccc [t/c]ttcccgccctggtgactgacggtctctgtgttgccttcaggacagc
22753	a	t	agcgaacagccttgaccctggctcagaatggcagttccttttttttt [a/t]ttattatttttttttttttttttttattgatcattcttgggtgtttcttgc
22945	g	a	gcagaacaaaatgaaaagtctcccatgtctacttctttctacacagacac [g/a]gcaaccatccgatttctcaatcttttccccacctttccccctttctatt
23032	g	a	ccccctttctattccacaaaaccgccattgtcatcatggcgcttctca [g/a]tgagctgttggtacacctcccagacggggtggtggccgggcagaggggc
23738	g	a	ggggctgcttggggtccatgagcacctttactcatatgtggggggcagaa [g/a]gctgtcccgtggtcagcagggccatgttggggcctcgggcagctgccgg
23952	t	g	tctgctcatgtcctgcactctgcagccaggcctgagaccaaggtacccc [t/g]ccgtggcctctgagtggggagtggtgctgggggcaggattcctgggcatgg
24123	a	g	gggcagtcaccttgctataaaccatggtgttctccactgtgtgggggcc [a/g]tgggggtctccccttagcacccctctcccggtccctgcagcacgctagg
24527	c	-	caggcatctctgagtggtgggctggaccgtcctctgtgggcctcagccagtgg [c/-]tgctgcagcaagggtggtggctccccacatatcactccttccctgcccct
24691	c	t	agggaaacagctgctgctgtcaccacaaattgtctctgtgctcctgccacc [c/t]gaatcccccaactgggcgactcagccgccacgagatggagcactctggcc
25015	g	a	gggcctgcagggtgggtgtgcactgggatgccttaggggttagcagctgcc [g/a]gaaggccttccctgtgcctgcctcccatggcgaatgtgacatctctgggct
25191	g	t	ctcgtccctccctgtggaagggggagcggtggccccagccctccgaggg [g/t]tcacgttatggcttctggtcactgccacactgtcctttgtgctggggaca

POSITION	Major	Minor	
30344	g	a	Intron
31170	a	g	Intron

16256	c	t	Intron
13376	a	t	Intron
12210	t	c	Intron
12072	g	c	Intron
11922	g	t	Intron
11903	-	a	Intron
10009	c	g	Intron
4519	c	t	Intron
4181	a	g	Intron
20952	a	c	Intron
20987	t	c	Intron
21620	g	a	Intron
21795	t	c	Intron
22753	a	t	Intron
22945	g	a	Intron
23032	g	a	Intron
23738	g	a	Intron
23952	t	g	Intron

24123	a	g	Intron
24527	c	-	Intron
24691	c	t	Intron
25015	g	a	Intron
25191	g	t	Intron

## Gene structure of Splice Form 2:

protein 1 MPSSVTALGQARSSGPGMAPSACCCSPAALQRRLPILAWLPSYSLOWLK  
 MPSSVTALGQARSSGPGMAPSACCCSPAALQRRLPILAWLPSYSLOWLK  
 MPSSVTALGQARSSGPGMAPSACCCSPAALQRRLPILAWLPSYSLOWLK  
 genomic 687 acttgagcgcgattgagcagttttcgccaaccacgtccattctca  
 tccctcctgacgcgcgtccgcgggccccctaggtcttcgtcgactagta  
 gttgggggtgcgtccgcgcgccccctgcgggggcggggccccggggg

protein 50 MDFVAGLSVGLTAIPQALAYAEEVAGLPPQ  
 MDFVAGLSVGLTAIPQALAYAEEVAGLPPQ  
 MDFVAGLSVGLTAIPQALAYAEEVAGLPPQ  
 genomic 834 agtgggctggcagaccgcgtgggggccccGTGAGGC Intron 1  
 tattcgtctgtcctcactcacatcggtcca<0-----[921 : 1781]  
 gtcccccatcctctcgggcttagtacgcg

protein 79 YGLYSAFMGCFVYFFLGTSRDVLGPTAIMSLLVSFYTFHEPAYAV  
 YGLYSAFMGCFVYFFLGTSRDVLGPTAIMSLLVSFYTFHEPAYAV  
 YGLYSAFMGCFVYFFLGTSRDVLGPTAIMSLLVSFYTFHEPAYAV  
 genomic 1779 CAGtgcttgtagttgtttcgatcgagcagcaatccgtttatcgcggtg  
 -0>agtaccttggttatttgccgatctgcccttctttctactaaccact  
 tccctccgcccgtccgcccgtgtgccctgcgcgccccctgccttg

protein 125 LLAFLSGCIQLAMGVRLRLAHISPHPLGLGGAGTSSMSPLGWPGFLLDFI  
 LLAFLSGCIQLAMGVRLRL P G L L F  
 LLAFLSGCIQLAMGVRLRLGEALPSCQG-----DALDLSICLFAFQ  
 genomic 1920 ccgtctgtaccgagcctggccttcg ggcgcaatttgtc  
 ttcttcggtatctgttggtgactccgag actatgtgttcta  
 ggccgccccggcgcggtgtgtattcag tccccctcgata

protein 174 SYPVIK FTSAAAVTIGFGQIK  
 YP ++ FTSAAAVTIGFGQIK  
 VYPRVR G:V[gtc] FTSAAAVTIGFGQIK  
 genomic 2040 gtccgcGTGTGTGCG Intron 2 AAGCtatggggaagtgcaa  
 tacgtg <2-----[2060 : 2347]-2> tccccctctgtgata  
 ctctgt cctttccccctagcg

protein 196 NLLGLQNI PRPFFLQVYHTFLRIAET  
 NLLGLQNI PRPFFLQVYHTFLRIAET  
 NLLGLQNI PRPFFLQVYHTFLRIAET  
 genomic 2394 GTAGGCA Intron 3 CAGaccgccaacacttccgtcatcaagga  
 <0-----[2394 : 4901]-0>attgtaatcgctttataacttgtcac

cggaagcccggccgggcccccgtagc

protein 222 VGDAVLGLVCM L L L L L V L K L M R D H  
VGDAVLGLVCM L L L L L V L K L M R D H  
R:R[agg] VGDAVLGLVCM L L L L L V L K L M R D H  
genomic 4980 AGGTACCCC Intron 4 CAGGggggggcgctaccccgcacacgc  
<2-----[4982 : 6882]-2> tgacttggtgtttttttattgaa  
atccccgggcccggggggggggggcc

protein 246 VPPVHPEMPPGVRLSRGLVWAATT  
VPPVHPEMPPGVRLSRGLVWAATT  
VPPVHPEMPPGVRLSRGLVWAATT A:A[gct]  
genomic 6953 gccgccgaccggccacgcgtggaaGGTGAGGG Intron 5  
tcctacatccgtgtgggttgcccc <1-----[7026 :10803]  
gtccccggccttgccctggcgtcga

protein 270 RNALVVSFAALVAYSFEVTGYQPFILTGETAEGLPVRI P P F S V  
RNALVVSFAALVAYSFEVTGYQPFILTGETAEGLPVRI P P F S V  
RNALVVSFAALVAYSFEVTGYQPFILTGETAEGLPVRI P P F S V  
genomic 10801 TAGCTcagcggttgccggtttggagtcctacaggagggcccgccaccttg  
-1> gactttctccttcactatcgaactttcgaccagtcctgtcctct  
cccggcccacgtgcccggtacgtccaaggatggctacgcgccag

protein 315 TTANGTISFTEMVQ DMGAGLAVVPLM  
TTANGTISFTEMVQ DMGAGLAVVPLM  
TTANGTISFTEMVQ DMGAGLAVVPLM  
genomic 10938 aagagaattagagcGTGGGCG Intron 6 CAGgagggcgggcca  
cccagctctcatta<0-----[10980:11397]-0>atgcgtcttctt  
caccggccccgggg cgacggcgggcgg

protein 341 GLLESIAVAKAF SQNNYRIDANQ  
GLLESIAVAKAF SQNNYRIDANQ  
GLLESIAVAKAF A:A[gca] SQNNYRIDANQ  
genomic 11434 gccgaaggaggtGGTAAGAC Intron 7 TAGCATcaatcaggac  
gttagtctcact <1-----[11471:14777]-1> caaaagtacaa  
ccggctggcacc tgttccctccg

protein 365 ELLAI LTNMLGSLVSSYPVTGSF  
ELLAI LTNMLGSLVSSYPVTGSF  
ELLAI G:G[ggg] LTNMLGSLVSSYPVTGSF  
genomic 14813 gccgaGGTAAGAC Intron 8 CAGGTcaaatgtcgtttcgagat  
attct <1-----[14829:15487]-1> tcattgcttcactcggg  
gggcc cccggccccccgcacct

protein 389 G TAVNAQSGVCTPAGGLVT  
G TAVNAQSGVCTPAGGLVT  
G TAVNAQSGVCTPAGGLVT  
R:R[cgg] TAVNAQSGVCTPAGGLVT  
genomic 15544 gCGGTGAGTG Intron 9 CAGGaggagctggtacgggcga  
g <2-----[15549:16445]-2> cctacacgtgcccggttc  
a acgctggggccggcggg

protein 409 VLVL LSLDYLTSLFYIIPKSALA  
VLVL LSLDYLTSLFYIIPKSALA  
G:G[gga] VLVL LSLDYLTSLFYIIPKSALA  
genomic 16501 GGTAAGGC Intron 10 CAGGAgcgccctcgtcatctttacatgcg  
<1-----[16502:16800]-1> tttttctaataccttaatacacctc  
gggggtgccgcagcccccgctcgt

26/27

intron	7026	10803
cds	10804	10979
intron	10980	11397
cds	11398	11470
intron	11471	14777
cds	14778	14828
intron	14829	15487
cds	15488	15548
intron	15549	16445
cds	16446	16501
intron	16502	16800
cds	16801	16941
intron	16942	18422
cds	18423	18550
intron	18551	18867
cds	18868	18967
intron	18968	19446
cds	19447	19580
intron	19581	21620
cds	21621	21693
intron	21694	22916
cds	22917	23005

## SEQUENCE LISTING

&lt;110&gt; APPLERA CORPORATION

<120> ISOLATED HUMAN TRANSPORTER PROTEINS,  
NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS,  
AND USES THEREOF

&lt;130&gt; CL000746PCT

&lt;140&gt; TO BE ASSIGNED

&lt;141&gt; 2001-10-26

&lt;150&gt; 09/765,344

&lt;15&gt; 2001-01-22

&lt;160&gt; 7

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 2919

&lt;212&gt; DNA

&lt;213&gt; Human

&lt;400&gt; 1

aacagcacga gggcggaacc agctgtggcg acgccaggag accccaagct gcatcgccga 60  
gtggaagcaa ctagaactcc agggctgtga aagccacagg tgggggctga gcgaggcggtg 120  
gcctcaggag cggaggagacc cccactctc cctcgagcgc cgcagtccac cgtagcgggt 180  
ggagcccgcc ttggtgcgca gttgaaaac ctggagccc cgctggatct cctggctgcc 240  
accgcacccc cccgccagcc tacgcccac cgtagagatg ccttctcgg tgacggcgct 300  
gggtcaggcc aggtcctctg gccccgggat ggcccagagc gcctgtgtgt gctcccctgc 360  
ggccctgcag aggagggtgc ccatctggc gtggctgccc agctactccc tcagtggtgt 420  
gaagatggat ttctgcgcc gctctcagt tggcctcact gccattcccc aggcgtggc 480  
ctatgtcga gttgctggac tccgcccc gtagtgccct tactctgcct tcatgggctg 540  
cttctgtat ttctctcgg gcacctccc ggaatgtgact ctgggcccc cgcattat 600  
gtccctcctg gtctcctct acacctcca tgagcccgcc tacgtgtgc tgctggcctt 660  
cctgtccggc tgcattcagc tggccatggg ggtctcgct ttgggggtcc tgctggactt 720  
catttcciac cccgtcatta aaggcttcac ctctgtgtgt gccgtcacca tcggcttgg 780

acagatcaag aacctgctgg gactacagaa catccccagg ccgttctcc tgcagggtga 840  
 ccacaccttc ctacaggattg cagagaccag ggtagggtgac gccgtcctgg ggctgggtctg 900  
 catgctgctg ctgctgggic tgaagctgat gcgggaccac gtgcctccc tccaccccga 960  
 gatgccccct ggtgtgcggc tcagccgtgg gctggtctgg gctgccacga cagctcgcaa 1020  
 cgccctgggt gtcctctcg cagccctggt tgcgtactcc ttcgagggtga ctggatacca 1080  
 gcctttcalt ctaacagggg agacagctga ggggtccct ccagtccgga tcccgcctt 1140  
 ctacgtgacc acagccaacg ggacgatctc ctacaccgag atggtgcagg acatgggagc 1200  
 cgggtggcc gtggtgcccc tgatggcct cctggagagc attgcggtgg ccaaagcctt 1260  
 cgcatctcag aataattacc gcatcgatgc caaccaggag ctgctggcca tcggctcac 1320  
 caacatgttg ggctccctcg tctctcta cccggtcaca ggcagcttg gacggacagc 1380  
 cgtgaacgct cagtcggggg tgtgcacccc gccggggggc ctggtgacgg gagtgtgtgt 1440  
 gctgtgtct ctggactacc tgacctact gttctactac atcccaagt ctgccctggc 1500  
 tgccgtatc atcatggcg tggccccgt gtcgacacc aagatctca ggacgtctg 1560  
 gcgtgttaag aggtggacc tgctgcccc gtgcgtgacc ttctgtgtgt gctctggga 1620  
 ggtgcagtac ggcattctgg ccggggccct ggtgtctctg ctcatgtcc tgcactgic 1680  
 agccaggcct gagaccaagg tgcagaggg gccggtctg gtctgcagc cggccagcg 1740  
 cctgtcttc cctgcatgg aggtctgctg ggaggagatc ctaagccgg ccttgaagt 1800  
 gtcccccca cgctgcctgg tctggagtg caccatgtc tgcagcatc actacactgt 1860  
 ggtgctggga ctggcgagc tctccagga ctccagaag caggggctg cctggcctt 1920  
 tggggcctg caggtcccc ttctcgtgt cctgtgtcc gctgacctg aggggtcca 1980  
 gtacttctt accctggaag aagcagagaa gcacctgagg caggagccag ggaccagcc 2040  
 ctacaacatc agagaagact ccatctgga ccaaaggtt gccctgtca aggcataatg 2100  
 gggccaccg tgggcatcca cagttgcag ggtgtccg aaggttctg tcatgtgat 2160  
 tggatgtgg atgccctg atagacatgc tggctggct gagaacccc tgagcaggta 2220  
 acccaggga gagaaggaag ccaggcctgg aggtccacg cagtgggagt ggggtcact 2280  
 ggctctctg gggatgactg gaaaatgacc tgcgtgtgt tccctggcat gacctctt 2340  
 ggaagagtgg ttggagaga gcctctaga atgacagact gtgcaggaa gcaggggcag 2400  
 ggggtccag ccgggtgtg gcgaggatc ctggggctgg cagcacctc ccggtcacc 2460  
 agtccacct gcgggggagg gacggggcag gcaggagtct gggaggcgg tccgtctc 2520  
 ttgtctcgg catctgtct cccgagaga aaaccaaggt gtgtcaaat acgtcaagt 2580  
 tctatttaa aataatttg tgtttctaa atggaaaaa tgatagctt ggtgatttg 2640  
 taaaagtc ataatgttat tgaaaaaa acaggaacc accctcacc ctgtccact 2700  
 gggatcatc tccagacccc tcccaaaca tgcatagtc cctgtccgtc agtgttgga 2760  
 tgaatglla cagttctaca taaatggat cattttatc atggtgtct ggaaccaca 2820  
 ttttcatgc agtcattgc agtgaattat ttattgtat aataaatagc attagaatac 2880  
 aagaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2919

&lt;210&gt; 2

&lt;211&gt; 2955

&lt;212&gt; DNA



&lt;213&gt; Human

&lt;400&gt; 2

gctgtggcga cgccaggaga cccaagctg catcgccgag tggaagcaac tagaactcca 60  
 gggctgtgaa agccacaggt gggggctgag cgaggcgtgg cctcaggagc ggaggacccc 120  
 cccactctcc ctgagcgcc gcagtccacc gtagcgggtg gagcccgctt tggcgcgag 180  
 ttgaaaacc tcggagcccc gctggatctc ctggctgcca ccgcacccc ccgccagcct 240  
 acgccccacc gtgagatgc ctcttcggt gacggcgctg ggtcaggcca ggtcctctgg 300  
 ccccgggatg gccccgagcg cctgtctgtg ctcccctgcg gccctgcaga ggaggctgcc 360  
 catctggcg tggctgcca gctactccct gcagtggctg aagatggatt tcgtcgccgg 420  
 cctctcagtt ggcctcactg ccattccca ggcgtggcc tatgtgaag tggctggact 480  
 cccgccccag tatggcctct actctgcctt catgggctgc ttcgtgtatt tctcctggg 540  
 caccctccgg gatgtgactc tgggccccac cgccattatg tccctcctgg tctcctcta 600  
 caccttccat gagcccgctt acgtgtgct gctggccttc ctgtccggtt gcattccagt 660  
 ggccatgggg gtctgcgtt tggctacat ctcccctcat cctctgggac tgggtggagc 720  
 cgggaccagc tcgatgtccc ctctggctg gccagggttc ctgtggact tcatttcta 780  
 ccccgctait aaaggcttca cctctgtgc tgcgtcacc atcggtttg gacagatcaa 840  
 gaacctgtg ggactacaga acatccccag gccgttctc ctgcagggtt accacacctt 900  
 cctcaggatt gcagagacca gggtaggtga cgccgtctg gggctggtt gcattgtct 960  
 gctgtgtgtg ctgaagctga tgcgggacca cgtgcctccc gtccacccc agatgcccc 1020  
 tgggtgcgg ctacccgtg ggtgtgtgtg ggtgccacg acagctcgca acgcccgtgt 1080  
 ggtctcttc gcagccctg ttcgtactc ctccagggtg actggatacc agccttcat 1140  
 cctaacaggg gagacagctg aggggtctcc tccagtccg atccgcctt tctcagtac 1200  
 cacagccaac gggacgatct cctcaccga gatgtgag gacatgggag ccgggctggc 1260  
 cgtgtgtccc ctgatggcc tctggagag caltgcgggt gccaaagcct tcgcatctca 1320  
 gaataattac cgcatcgtg ccaaccagga gctgtggcc atcggttca ccaacatgtt 1380  
 gggctccctc gtctctct acccggtcac aggcagctt ggacggacag ccgtgaacgc 1440  
 tcagtcgggg gtgtgacccc cggcggggg cctggtgacg ggagtgtgtg tctgtgtc 1500  
 tctggaclac ctgacctac tgttacta catcccaag tctgccctg ctgccgtcat 1560  
 catcatggcc gtggccccgc tttcgacac caagatctt aggcagctct ggcgtgttaa 1620  
 gaggctggac ctgtgcccc tgtgggtgac ctctctgtg tcttctggg aggtgcagta 1680  
 cggcatctg gccggggccc tgggtctct gctatgtc ctgcactct cagccaggcc 1740  
 tgagaccaag gtgtcagagg ggcgggttct ggtcctgcag ccggccagcg gcctgtcct 1800  
 ccctgcatg gaggctctg gggaggagat cctaagccg gccctggaag tgtcccgcc 1860  
 acgtgcctg gtctggagt gcacccatgt ctgcagcat gactacact tgggtgtgg 1920  
 actggcgag ctctccagg acttcagaa gcaggggct gccctggcct ttgtggcct 1980  
 gcaggtcccc gttctccgt tctgtgtc cgtgacctg aagggttcc agtacttct 2040  
 taccctggaa gaagcagaga agcacctgag gcaggagcca gggaccagc cctacaacat 2100  
 cagagaagac tccattctg accaaaaggt tgcctgtc aaggcataat gggccaccc 2160  
 gtgggcatcc acagtttga ggggttcc gaaggttct gtcactgtga ttgatgtc 2220

gatgccgct gatagacatg ctggcctggc tgagaaaccc ctgagcaggt aaccagga 2280  
 agagaaggaa gccaggcctg gaggtccacg gcagtgggag tggggctcac tggcttctg 2340  
 tgggatgact ggaaatgac ctgctgctg ttccctggca tgacctctt tgaagagtg 2400  
 gtttgagag agcctctag aatgacagac tgtgcgagga agcaggggca ggggttcca 2460  
 gcccgggctg tgcgaggcat cctggggctg gcagcacctt cccggctcac cagtgcacc 2520  
 tgcgggggag ggacggggca ggcaggagtc tgggaggcgg gtccgctcct ctgtctgcg 2580  
 gcatctgtc tctccgagag aaaaccaagg tgtgtcaat gacgtcaagt ctctattaa 2640  
 aaataattt gtgtttcta aatggaaaaa gtgatagctt tgggtattt gtaaaagtca 2700  
 taaatgctta ttgtaaaaaa tacaggaaac caccctcac cctgtccact tgggtgatca 2760  
 ttccagacc ctcccaaac atgcatatgt acctgtcgt cagtgtgtg atgtatgtt 2820  
 acagtctac ataaatggga tcatttata catgtgtc tggaaccac attttcatg 2880  
 cagtcattg cagtgaatta ttattgtga taataaatag cattagaata caaaaaaaaa 2940  
 aaaaaaaaa aaaaa 2955

<210> 3

<211> 606

<212> PRT

<213> Human

<400> 3

Met Pro Ser Ser Val Thr Ala Leu Gly Gln Ala Arg Ser Ser Gly Pro  
 1 5 10 15  
 Gly Met Ala Pro Ser Ala Cys Cys Cys Ser Pro Ala Ala Leu Gln Arg  
 20 25 30  
 Arg Leu Pro Ile Leu Ala Trp Leu Pro Ser Tyr Ser Leu Gln Trp Leu  
 35 40 45  
 Lys Met Asp Phe Val Ala Gly Leu Ser Val Gly Leu Thr Ala Ile Pro  
 50 55 60  
 Gln Ala Leu Ala Tyr Ala Glu Val Ala Gly Leu Pro Pro Gln Tyr Gly  
 65 70 75 80  
 Leu Tyr Ser Ala Phe Met Gly Cys Phe Val Tyr Phe Phe Leu Gly Thr  
 85 90 95  
 Ser Arg Asp Val Thr Leu Gly Pro Thr Ala Ile Met Ser Leu Leu Val  
 100 105 110  
 Ser Phe Tyr Thr Phe His Glu Pro Ala Tyr Ala Val Leu Leu Ala Phe  
 115 120 125  
 Leu Ser Gly Cys Ile Gln Leu Ala Met Gly Val Leu Arg Leu Gly Phe  
 130 135 140  
 Leu Leu Asp Phe Ile Ser Tyr Pro Val Ile Lys Gly Phe Thr Ser Ala  
 145 150 155 160

Ala Ala Val Thr Ile Gly Phe Gly Gln Ile Lys Asn Leu Leu Gly Leu  
 165 170 175  
 Gln Asn Ile Pro Arg Pro Phe Phe Leu Gln Val Tyr His Thr Phe Leu  
 180 185 190  
 Arg Ile Ala Glu Thr Arg Val Gly Asp Ala Val Leu Gly Leu Val Cys  
 195 200 205  
 Met Leu Leu Leu Val Leu Lys Leu Met Arg Asp His Val Pro Pro  
 210 215 220  
 Val His Pro Glu Met Pro Pro Gly Val Arg Leu Ser Arg Gly Leu Val  
 225 230 235 240  
 Trp Ala Ala Thr Thr Ala Arg Asn Ala Leu Val Val Ser Phe Ala Ala  
 245 250 255  
 Leu Val Ala Tyr Ser Phe Glu Val Thr Gly Tyr Gln Pro Phe Ile Leu  
 260 265 270  
 Thr Gly Glu Thr Ala Glu Gly Leu Pro Pro Val Arg Ile Pro Pro Phe  
 275 280 285  
 Ser Val Thr Thr Ala Asn Gly Thr Ile Ser Phe Thr Glu Met Val Gln  
 290 295 300  
 Asp Met Gly Ala Gly Leu Ala Val Val Pro Leu Met Gly Leu Leu Glu  
 305 310 315 320  
 Ser Ile Ala Val Ala Lys Ala Phe Ala Ser Gln Asn Asn Tyr Arg Ile  
 325 330 335  
 Asp Ala Asn Gln Glu Leu Leu Ala Ile Gly Leu Thr Asn Met Leu Gly  
 340 345 350  
 Ser Leu Val Ser Ser Tyr Pro Val Thr Gly Ser Phe Gly Arg Thr Ala  
 355 360 365  
 Val Asn Ala Gln Ser Gly Val Cys Thr Pro Ala Gly Gly Leu Val Thr  
 370 375 380  
 Gly Val Leu Val Leu Leu Ser Leu Asp Tyr Leu Thr Ser Leu Phe Tyr  
 385 390 395 400  
 Tyr Ile Pro Lys Ser Ala Leu Ala Ala Val Ile Ile Met Ala Val Ala  
 405 410 415  
 Pro Leu Phe Asp Thr Lys Ile Phe Arg Thr Leu Trp Arg Val Lys Arg  
 420 425 430  
 Leu Asp Leu Leu Pro Leu Cys Val Thr Phe Leu Leu Cys Phe Trp Glu  
 435 440 445  
 Val Gln Tyr Gly Ile Leu Ala Gly Ala Leu Val Ser Leu Leu Met Leu  
 450 455 460  
 Leu His Ser Ala Ala Arg Pro Glu Thr Lys Val Ser Glu Gly Pro Val  
 465 470 475 480

Leu Val Leu Gln Pro Ala Ser Gly Leu Ser Phe Pro Ala Met Glu Ala  
 485 490 495  
 Leu Arg Glu Glu Ile Leu Ser Arg Ala Leu Glu Val Ser Pro Pro Arg  
 500 505 510  
 Cys Leu Val Leu Glu Cys Thr His Val Cys Ser Ile Asp Tyr Thr Val  
 515 520 525  
 Val Leu Gly Leu Gly Glu Leu Leu Gln Asp Phe Gln Lys Gln Gly Val  
 530 535 540  
 Ala Leu Ala Phe Val Gly Leu Gln Val Pro Val Leu Arg Val Leu Leu  
 545 550 555 560  
 Ser Ala Asp Leu Lys Gly Phe Gln Tyr Phe Ser Thr Leu Glu Glu Ala  
 565 570 575  
 Glu Lys His Leu Arg Gln Glu Pro Gly Thr Gln Pro Tyr Asn Ile Arg  
 580 585 590  
 Glu Asp Ser Ile Leu Asp Gln Lys Val Ala Leu Leu Lys Ala  
 595 600 605

&lt;210&gt; 4

&lt;211&gt; 630

&lt;212&gt; PRT

&lt;213&gt; Human

&lt;400&gt; 4

Met Pro Ser Ser Val Thr Ala Leu Gly Gln Ala Arg Ser Ser Gly Pro  
 1 5 10 15  
 Gly Met Ala Pro Ser Ala Cys Cys Cys Ser Pro Ala Ala Leu Gln Arg  
 20 25 30  
 Arg Leu Pro Ile Leu Ala Trp Leu Pro Ser Tyr Ser Leu Gln Trp Leu  
 35 40 45  
 Lys Met Asp Phe Val Ala Gly Leu Ser Val Gly Leu Thr Ala Ile Pro  
 50 55 60  
 Gln Ala Leu Ala Tyr Ala Glu Val Ala Gly Leu Pro Pro Gln Tyr Gly  
 65 70 75 80  
 Leu Tyr Ser Ala Phe Met Gly Cys Phe Val Tyr Phe Phe Leu Gly Thr  
 85 90 95  
 Ser Arg Asp Val Thr Leu Gly Pro Thr Ala Ile Met Ser Leu Leu Val  
 100 105 110  
 Ser Phe Tyr Thr Phe His Glu Pro Ala Tyr Ala Val Leu Leu Ala Phe  
 115 120 125

Leu Ser Gly Cys Ile Gln Leu Ala Met Gly Val Leu Arg Leu Ala His  
 130 135 140  
 Ile Ser Pro His Pro Leu Gly Leu Gly Gly Ala Gly Thr Ser Ser Met  
 145 150 155 160  
 Ser Pro Leu Gly Trp Pro Gly Phe Leu Leu Asp Phe Ile Ser Tyr Pro  
 165 170 175  
 Val Ile Lys Gly Phe Thr Ser Ala Ala Ala Val Thr Ile Gly Phe Gly  
 180 185 190  
 Gln Ile Lys Asn Leu Leu Gly Leu Gln Asn Ile Pro Arg Pro Phe Phe  
 195 200 205  
 Leu Gln Val Tyr His Thr Phe Leu Arg Ile Ala Glu Thr Arg Val Gly  
 210 215 220  
 Asp Ala Val Leu Gly Leu Val Cys Met Leu Leu Leu Leu Val Leu Lys  
 225 230 235 240  
 Leu Met Arg Asp His Val Pro Pro Val His Pro Glu Met Pro Pro Gly  
 245 250 255  
 Val Arg Leu Ser Arg Gly Leu Val Trp Ala Ala Thr Thr Ala Arg Asn  
 260 265 270  
 Ala Leu Val Val Ser Phe Ala Ala Leu Val Ala Tyr Ser Phe Glu Val  
 275 280 285  
 Thr Gly Tyr Gln Pro Phe Ile Leu Thr Gly Glu Thr Ala Glu Gly Leu  
 290 295 300  
 Pro Pro Val Arg Ile Pro Pro Phe Ser Val Thr Thr Ala Asn Gly Thr  
 305 310 315 320  
 Ile Ser Phe Thr Glu Met Val Gln Asp Met Gly Ala Gly Leu Ala Val  
 325 330 335  
 Val Pro Leu Met Gly Leu Leu Glu Ser Ile Ala Val Ala Lys Ala Phe  
 340 345 350  
 Ala Ser Gln Asn Asn Tyr Arg Ile Asp Ala Asn Gln Glu Leu Leu Ala  
 355 360 365  
 Ile Gly Leu Thr Asn Met Leu Gly Ser Leu Val Ser Ser Tyr Pro Val  
 370 375 380  
 Thr Gly Ser Phe Gly Arg Thr Ala Val Asn Ala Gln Ser Gly Val Cys  
 385 390 395 400  
 Thr Pro Ala Gly Gly Leu Val Thr Gly Val Leu Val Leu Leu Ser Leu  
 405 410 415  
 Asp Tyr Leu Thr Ser Leu Phe Tyr Tyr Ile Pro Lys Ser Ala Leu Ala  
 420 425 430  
 Ala Val Ile Ile Met Ala Val Ala Pro Leu Phe Asp Thr Lys Ile Phe  
 435 440 445

Arg Thr Leu Trp Arg Val Lys Arg Leu Asp Leu Leu Pro Leu Trp Val  
 450                      455                      460  
 Thr Phe Leu Leu Cys Phe Trp Glu Val Gln Tyr Gly Ile Leu Ala Gly  
 465                      470                      475                      480  
 Ala Leu Val Ser Leu Leu Met Leu Leu His Ser Ala Ala Arg Pro Glu  
 485                      490                      495  
 Thr Lys Val Ser Glu Gly Pro Val Leu Val Leu Gln Pro Ala Ser Gly  
 500                      505                      510  
 Leu Ser Phe Pro Ala Met Glu Ala Leu Arg Glu Glu Ile Leu Ser Arg  
 515                      520                      525  
 Ala Leu Glu Val Ser Pro Pro Arg Cys Leu Val Leu Glu Cys Thr His  
 530                      535                      540  
 Val Cys Ser Ile Asp Tyr Thr Val Val Leu Gly Leu Gly Glu Leu Leu  
 545                      550                      555                      560  
 Gln Asp Phe Gln Lys Gln Gly Val Ala Leu Ala Phe Val Gly Leu Gln  
 565                      570                      575  
 Val Pro Val Leu Arg Val Leu Leu Ser Ala Asp Leu Lys Gly Phe Gln  
 580                      585                      590  
 Tyr Phe Ser Thr Leu Glu Glu Ala Glu Lys His Leu Arg Gln Glu Pro  
 595                      600                      605  
 Gly Thr Gln Pro Tyr Asn Ile Arg Glu Asp Ser Ile Leu Asp Gln Lys  
 610                      615                      620  
 Val Ala Leu Leu Lys Ala  
 625                      630

&lt;210&gt; 5

&lt;211&gt; 31766

&lt;212&gt; DNA

&lt;213&gt; Human

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(31766)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 5

ggtccccgcg gccctcggcc ttgctcgggg ccaagggacc gcggacggtc aggtggcgca 60  
 gggctcctc cggagacccc aggatccgga gccagcggcc ttgtgggcag gggccggggg 120  
 caggggagtg gattttgccc ggagcggagc aggccggggg cagtgggggg ctgggggtga 180

gggtggctgg ctctgcgcgc gggcgccggg gccctggaag atgctgcgca cctgaattaa 240  
ccggggcgct ctgatgtcct cccagaagca actagaactc cagggctgtg aaagccacag 300  
gtgggggctg agcgaggctt ggcctcagga gcggaggacc ccccccccc cccctcgag 360  
cgccgcagtc caccgtagcg ggtggagccc gccttggtgc gcagttgaa aacctcgaa 420  
gccccgctgg atctcctggc tgccaccgc acccccgcc agctacggtg cgccgcggg 480  
ccagcttct ctctgcgtg ctccccgta aattccctgg ggagacggaa aaaaaggcaa 540  
aggaagtcgg ttctccaggg gccagaagtg ttgagcctaa ttagtctca gacttctca 600  
tgaggaatcg ctatcagtt tctatctgg gagagttgag gatggaggga cagaaggcac 660  
ccaggatttg cacggggggg gattcaggga gagaggggta tgaggacgg ggtgggcctt 720  
ccagcttgg cccagtcctc atctgcaca cattgttgc ttctcttag agccgttcgc 780  
ccccctggg aggggagacc catagtacc tctctgaca ccgcccacc ctgaccagt 840  
ttgccgggtt ctcaaaggc cagctctga ctgctggtc gtgtacctg cccccccag 900  
ccccaccga gagatgcctt ctgggtgac ggcgtgggt caggccaggt cctctggccc 960  
cggaatggcc ccgagcgct gctgctgct cctgcggcc ctgcagagga ggctgccat 1020  
cctggcgtgg ctgccagct actccctga gtggctgaag atggattcg tcgccgcct 1080  
ctcagttggc ctactgcca ttcccaggc gctggcctat gctgaagtgg ctggactccc 1140  
gccccaggtg aggcgtctga cctgctgcc agccatact cagaaacagt gcagaatca 1200  
cagatcaat cccagacacc atcagcgatt ccaggttcc agccccggg cccaaggaa 1260  
ccttggtt acagtgtg acgcagatt tctctggcc gaccaggct cctatgcctg 1320  
ttgttacac acagacactg agctggtat ggaggggcca gcgagatgac tcatggaggc 1380  
ctcaggagt caagaccagc ccgacaaaaa tggtaaaacc ccgtctccac taataataca 1440  
aaaattaggc tgggtgcgtt ggctcaagcc tgtaatcca gcacttggg aggcggaggc 1500  
aggcggatcg caaggtcagg agatggagac catcctggct aacagggtga aacctgtct 1560  
ctactaaaaa taacaaaaa tagccagggtg tggtgccggg tgctgtagt cccagctacc 1620  
tgaggaggct aggcaggaga atggcgtgaa ccggggaggc ggagcttga gtgagccgag 1680  
actgtgccac tgcctccag cctgggcgac agagcgagac tcatctcaa aaaaaaaaaa 1740  
aaaagaatgc ttctcagac ttggacacag cacacgggcc tgtaccgacc cctctgcctg 1800  
gctgtctga cctgaggcc ccagttgagt gctgctaaa aagtggcctc ctgatcactg 1860  
caggtccacc cacagggcag ggcgtgcac cttaacctg ggcctggaca cagctgacac 1920  
ccacacatcc cgagcttga cagcacact agggagctgg tggatggcc tcggcctct 1980  
gagtgtcac caccctct cccacagta tggcctctac tctgcctca tgggtgctt 2040  
cgtgtattc ttctgggca cctccggga tgtgactct ggccccacc ccattatgc 2100  
cctcctggtc tcttctaca ccttcatga gccgcctac gctgtgtgc tggccttct 2160  
gtccggtgc atccagctg ccatgggggt cctgcgttg ggtgaggctc taccttctg 2220  
ccaaggggat gccctgacc tcagcattg ctgtttgca ttcaagtct atccccgtg 2280  
gcgtgtgtg gcgtgtgtg ggggtgtgg tatgtatgt tgtgtgtga ggtgggtgg 2340  
tggtgaggg ggtggggcac ttgctcct agtctactat ttactgatt agaggccagg 2400  
acattggaga aagtgacctg tggctcagac cccatatgcc ccccccccc nnnnnnnnn 2460  
nnnnnnnnn nnnnnnnnn nnnnnnnnn nntcattc ctacccgctc attaaagct 2520  
cacctctgt gctgccgta ccatcggtt tggacagatc aaggtaggca cggcgccac 2580

ccagggcact gctcttggc cactgctcgt tggcacaggg atggcgggag caggactgag 2640  
 gccagtcctg atccctgtgg ccagtgagcg tctgtctgt tcagattgc tccatgggt 2700  
 caagaagcac gcggtgctct catgggtccc ctgttaataa aatgaccclc ctgggagggg 2760  
 gtacacgtga tgggtggatt tcacagcggg taactgggg gccggttaatt ccatccccct 2820  
 gctcttggc gagttccgt gccagtgtgc ttggctggct ctgtacgtg gctctgtct 2880  
 cctgcactg ggcacaccca gcaggcccca ccagtcatga gcatgtcgt agaatttcta 2940  
 taggcaaatt atttcccat gccaattag ttagatgggt ttgtttgtg cttttttc 3000  
 tggttctgt ttaatatatt taaatgcc a tgcctttat atttctttt aaaacattta 3060  
 taataagtac taaaatcagt catcagctga ggggttaatt tattctgtt ttgctgggt 3120  
 glaagttcct cgagggtagg aactgtcct ctcccagcca tggccgcca gcallgggt 3180  
 ggtgcagtag ggggggtcct agtgggtgt gtgtggagt aagtggccc agttccacag 3240  
 atgggacct gcggccctca tggcagacta ggtcacatg ctgcctctg accctgtgc 3300  
 actgcagtc atcttccca agccaggccc tgtccaggc tggcctgaga cagtctccc 3360  
 tgatggaggt accatgagaa gaccaaggac aggagagtgt gtgtgagagt gtgtatgagt 3420  
 gtgtgtgtgt gtgaatgagt gagtgtgta gagtgtgagt gggttgagg gagtgtgtgt 3480  
 gtgcgtgtgt gagtgtgtgt gagtgtgggt gtgagtgagt gcgtgtgtgt gagagagtgt 3540  
 gtgagtgagt atgagtga ggtgtgaggt ggttagggagt gtgtgagtgt 3600  
 gcgcgcgcgt gtgtgagtgt atgagtgtga gagtgtgtgt gagtgtgaga gagtgggtgt 3660  
 ggggtgtagt gtgcgtgtga gtgtgagagt gtgagtgagt gagtgtgaga ataaagtaga 3720  
 cacttttgc actcttgcta cgtgcgaggc actgggcagg acacttcca tgttaattc 3780  
 tcaacacacc cgggagagta gatttatcat tattccatt tgagagagga gggaccaact 3840  
 taggtgtggg tgagcgtgat ttgtacgta tctacagca tcttcaggg ttagctggc 3900  
 aaatgctgt tcagggcatg gttggttct taatctggaa acatcattt tgggtcaag 3960  
 aatgtcttt ttaggatcc cagtgtgagt ggagagcggg gagtggagag tggagggcat 4020  
 ccttgttca tgcctcata ctggcaacc ctgccccgc ccagggactc tcagccatc 4080  
 tggggggagg ggcgtctcc tgacaggccc aggacagaag accctacccc aacagtcca 4140  
 gagtccgccc ccaggtatgt cctaacccca cccacctgt acgcagcagt ttagggcag 4200  
 ggtccttgag cctgtggcca tgggatccag ggcctcaact tccccctg actcctgtt 4260  
 ggtcagggc gatgtaaaga actaaagtgc aagctgtcc tgcactgtg atccagcta 4320  
 cgtgggaggc cgaggtggga ggtactgtg agccaggag ttgaatcca gcctgggcca 4380  
 cagagcaagc aagaccccca cccccaacc ccccgccgt ctctaaagaa aaaaaacta 4440  
 aactaaaaca caggatagag tgccttctc tggccaggac ctgagagctg gtagctgtgt 4500  
 gggaggctcc tactttgcc aggatcccc aagctggtt ctgaagccc ctgagagccc 4560  
 tccacatctg cacactacga agagatttt cctccgcag cagcggagct ggggggtggc 4620  
 ggggcaccta gtgagggaga catttcaag ccaatggcag caagggtc cactgcaggg 4680  
 ggcctggctg ctgagtggc tggagccaat ggggtgggg ccaggcagcc catggccct 4740  
 ggccatcaag ctgtagaacc tgtctgctg ctgtgggtc acctacattg tttttatga 4800  
 gaaaagtaat ttagagaaaa acatatcact gaccagtaa ttgtgaagta ctgtctcca 4860  
 cgagggtagc ttgatctcc tgcctaggg ggcgttggga gtgggcaggt ggacctga 4920  
 gccclaaaga tgggccccg gaataagggt tggagcagg gccgggggac actgtctcag 4980



ccctagggga ggtggcgagg gagctggga cagatggcct tggttggga gcatagcctc 5040  
tgatcagcat ctctgtgtt ggacagaacc tgctgggact acagaacatc cccaggccgt 5100  
tcttctgca ggtgtaccac accttctca ggattgcaga gaccaggtag cccgggctt 5160  
gttctccct cctataagga agtctctt tccacacctc ctctccggc cccacactca 5220  
gttccccac cctggtgac tgctaaaca ggggtccca gagcagccc atcagcagca 5280  
cctgcaagct ggcaagaaat gagacttca gtccttgca gacctgcct atcgagacca 5340  
gagglttaac aggaggccgg tggctgagc gcaggggaga gttcaagaag ctctgtctg 5400  
gggagctgga gcagggcaat gttctctct ccacaaagct ttctggggg acaggaatgg 5460  
ggagagtctg tglaaagaaa aagaaggcag tgtgtctcc ccaggctgtg attgttaag 5520  
gaggagaaca caggcggtgg ggagctaacc cagacagaac gcagggtgtg caggacggc 5580  
agggtgagct gtgatgagag atgaggagac ccaagcactt cctcagggtg gtttaagag 5640  
cctctgatgg cggacactg tggctcacac ctgtaatccc agcactttg gaagcccagg 5700  
tgggcagatc atcttaggtc aggagtcaa gaccagctg gtcaaatgg tgaactccg 5760  
tcttactaa aaatacaaaa attagtggg tgtgatggtg cgtgcctgta gtccagctg 5820  
ttcaggaggc tgaggcagga ggaattgtt aacctgtgag gcggagggtg cagtgtgctg 5880  
agatcacgcc actgcactcc agccgggtg atagagttag actccatctc cgcctgccc 5940  
tgccccccg aaaagaaaag cctctgatga ggggtacctc cctgccagac catccagcg 6000  
gaaggcagga tgctctcta cctctgtg gcgaagggg tggaggagga aatgatctag 6060  
gggagctata gagatggtg cccagtgtg tggcctggag ggagtgggg cgatcctgga 6120  
ccatctccc ctcccctg atctgtgccc gaagctccc agcagctgaa cccagctgga 6180  
agccagtggg ttctgtctc ttcagaggca ccagtgggca ggggtcggc agatggggga 6240  
gcagtatta cggagcctga ccagtcagag agcgtcacgg gctgtcatgt aaaaggggc 6300  
ttactgaccg ttagtctta agccttctag aatgtctaga gcagggtgtc cagtctttg 6360  
tcttccctg gccacacagg aagaattgtc ttaaaatata claatgatag ctgatgagct 6420  
acaaaaaat cattgaaaga actcataatg tttaagaaa gttacaaat ttgtgtggg 6480  
ccacattcaa agccatccca ggctgcacat ggcccacagg ctgtgggtt ggcagacttg 6540  
ttctagagcc acctgcagag agcatcagct gctcaggctg gtcccatc cctgtgtca 6600  
ctcaccagat cctgttcat agactgagt cccagagagc tacgggagt aaaatgtgag 6660  
cacgtcgtc tgcatcatc atgaccggg cctctccact gctgtcctg cccaagcctg 6720  
tgtacttact gaatttgaa ttgagtttg aagtagaaca gggatgcata ttgcttagga 6780  
gaaaacatct cagcctagtc tgttttgg cttgcagagt agcttctaa agttcctgag 6840  
cttagttaa gtgtctgta aatgggaaa atattgaaa tgctttatt cctgtgata 6900  
ctcactgtg tctgggacca gcagcctgg cattgggct cctgggagc ttctagtgc 6960  
ctctcagct aagcagcagc aggacgctg agtcagcagt aggcctggc cgagctgggt 7020  
gggtaccagt cctctgctg tccacagggt aggtgacgcc gtctggggc tggctgcat 7080  
gctgtctgt ctggtgtga agctgatgc ggaccagtg cctccgctc accccagat 7140  
gccccctgt gtgcggctca gccgtgggt ggtctgggt gccacgacag gtgaggggc 7200  
tctggctgac atcgtatga acctggctg caggtgggg tcaattggg agtcctagt 7260  
ccaccctagg gattctacg tcattggtct ggggtcact tgagcattgg gacatttaa 7320  
acaccacacc aaactctgg acatgtactt ttatttaat taattaatta attattttt 7380

gagacagagt ttgctctgt cgcccaggct ggagtacagt ggtgtatat tggctcactg 7440  
caacctccgc ctcccagggt catgtgattc tctgcctca gcctcctgag tagctggat 7500  
tacaggcgca tgcaccacg ctgactgat tttgtlalt atagtagaga cggggttca 7560  
ccatgtgac catactggtc tcaactcct gacctcagat gatctgcctg cctcgccctc 7620  
ccgaagtgt gggattacag gcatgagcca ccgccccgac ccaactcta gaacatttaa 7680  
aactctgagc caaactcggg gggttctgat gcagggtcag agctgagaac tgatcatgg 7740  
ccagctgtgg ccgttctcaa cctgaagcag tttgcccc ctggtgacat gtggcaatgt 7800  
tgggaaacat tttgttga taccaccggg cgaggcaggt gctggcaact agcgaggagg 7860  
ggccaggaat gccactaagc ccttgacggc gcacagcctt ccacagcaaa cagtatccg 7920  
gccccaaatg tcggcgggag cgagggtgag aaacacagtc tactggcagg tctggggag 7980  
acagaacaaa ttccggggag accaacaggc tctgttgaa ttgtgcagct ttatgggggt 8040  
acaacttcag cttcaggcgc ggagggtgca ggctgagcca caaccaacca ggggtcctt 8100  
gagaaggggg tggtagggc tgcgtcgtg aggagcagg ctgtcagcag gtgaaggacc 8160  
accttggggg aggtggaagc ctcttcag tgcattccgg atgggcatgg gtccctggc 8220  
tgagcaccgg gcagggatgc ccagagaaga aggcacacc aggaagtca ggaagcatag 8280  
gtactcca acccttcag agccgagggg acagattaag ctcaataa cccgagctg 8340  
gtctgtgt ttgggggctg caaatccaa cagccacaga agagagaggg tggctggatt 8400  
tagctctga gtcactctg cctgccagg ctggtctgt ctctgtgtc tctccctga 8460  
tctgttcag cagactgag gccgtgtga acactgagag tctgtgggg tgcaggat 8520  
gttgggtta gattggagat gtccagctc cagtcaccgc ctggcctgg ttagactg 8580  
caaagtagct tcaaagtc ctgagctta gtacagttg ccgtgaaata gtgaaatgc 8640  
ttgtatct ttgtatct ttgttttg ttttttg gagacgggt ctactcctg 8700  
ttgccaggc tggagtgcac tggcgtgat ccttggtca ctgcaacct cctcctggt 8760  
ttcaagcaat tattcccc ttaacctcc cgagtagcc tgggattac aggcgtgca 8820  
cccaccac ccgggcta tttgtatt ttgagttca agatggggg ttcaccattg 8880  
ttggccatg ctgggtctg aaactccnn nnnnnnnnn nnnnnnnnn nnnnnnnnn 8940  
nnnnnnnnn nnnnnnncc taacctga aaggcaggt ctccgagcc agaaaggtct 9000  
ttaagatg ttagacatg gtaatgcca gaaactcag aaacattta cagttaggg 9060  
ctgtgactc ctttcccc agtgtgtgta gattgtctt aggaactac taaaaagct 9120  
ttgatctt ataacagct tgagaatgag ttattatt ttccattt gtcagtagg 9180  
aaagttagcc tatgaaagat gaaaaataa acagaactc tcaaggccac acgactggga 9240  
agaggcagaa caaggactc acccgagtcg tctgtcttt ttctgaatg gtcgtctag 9300  
aagtgtgca atgtatgat ccttcaaag aaccagctt tggttcatt gatccctt 9360  
ttgcttgg ttctagtc attgattct gctgggct ttattttt ctttctta 9420  
ttatttca gtaactta tctttttt tttttt aaaaaaat tttttt 9480  
gagacggagt ttgctctg tccccaggg tggagtcaa tgggtgac tcagctact 9540  
gcaacctcg cctccgggt tcaagtatt ctctgcctc agcctccaa gtagctggga 9600  
ttacaggct atgccacat gcctggctaa tttttgtat tttaataga gacggggtt 9660  
caccatgtg gccaggctg tctgaactc ctgaactcag gtatccacc cgcctcagc 9720  
tccaaagt ctgggattac aggcgtgag cactgtgcc gccctttt ttacattct 9780

taaggaggaa gccctgggctc ttgatttgag acccttctc ttttlaata tgggtgtta 9840  
 aatgctgtaa atttactcc tagcattact ttgctgcat cccacagtt ttatatatat 9900  
 gctgtcttt aatttttatt cgggtctaaa tactttgtga ttccatttt gactctcct 9960  
 ttgacctgtg gattatttag aagtggatta tatgattct tctgtgaagc atgggagta 10020  
 tttagttcc aaatagtgg gggtttcca attagtact tgccattgat ttctaatta 10080  
 gttccattgt gataagaaaa cgtacttagt gattaaatc tgtttcctt tattgagagt 10140  
 tgtattatgg ccttgaatat ggtctatctt agtgaatatt ctgtgtgtgc ttgagaacat 10200  
 gttttctgcc attattgggt ggaatgtct gtaaatgca gttaggctg gttgggtatc 10260  
 tcaggctact atattctgc tgattttct tctactgtt ctgttgatta ttaaaagaag 10320  
 aatgtgaaa taactcagc tataaccatg acttgccta ttctcttc ttactacca 10380  
 gttattgtt catatattt gaagctctgt ttttagtat aaaaacatat agaattataa 10440  
 ttctcttg gaatgatact tttttctg ggcaaatc tctccctga aatctactt 10500  
 gatacaga gagctactcc agcttcatt agtgttagca tatcttcc catctttta 10560  
 cttttttt ttgagacag aatctctc tggtgccag gctggagtgc aatggcgtga 10620  
 tctggctca ctgcaacct cactccag gttcaagcaa ttgtgtgcc tcagcctccc 10680  
 aagtagctgg aattacaggc tgcaccacc aagccagct aattttgt ttttagtag 10740  
 caatgaggtt tcacatgtt ggccagcct gtcctcaatt ctggcctca agtgatccac 10800  
 ctgcttgc ctcctaaagt gctgggacac cacaccagc ctctatctt ttacttta 10860  
 acgtattaa ttgatttg ttctgtgt taatggtatg tttatggc agcatataga 10920  
 tgggattgt tttaatctc ataactctg cttttaatt ggaacttat gccattata 10980  
 tttaagtga ttattgct cttgactt agactacca tgtgtgtt tgttctgt 11040  
 ttgtctatt tgttctgt tcccttcc cctctctt acctactg gattatttt 11100  
 tatgattcca ttacatcc ttattggct gattagctg aacggctata tttagtttt 11160  
 gttgttgt ttctaagtga ctgcttaag gtttggtta aacatctta tctgtlaca 11220  
 gtataatta aactcaact tcatgtata taagaacct ataacaggcc aggcgcagt 11280  
 gctcacacct ataactctag cacttggga ggctgagga ggcagattgt ctgagctcag 11340  
 gagtcaaga ccagcctgg caacatggc aaacccatc tctacaaaa atacaaaaa 11400  
 ttaggcaggc atggtgtgc atgctgtaa tccagctac tgggaggct gaggtggag 11460  
 gattgctga acctggaagg cagagattgc agtgagcaa gatccgtcac tgaagctga 11520  
 aactcctgg ctcaagta tctctgcct cagcctctg agtagctgg tctacaggca 11580  
 tgcaccacca catctggcta atttttaa tttttata gaaacgggt atgtgtca 11640  
 ggctgatcaa aggttctgg cctcaagtga tgcctgcc tcagccaacc aaagtgtga 11700  
 gattgcagat gtgagccacc attccctga ggaacagct tagattatc cagtagtca 11760  
 ctgttctg tgcctta tctttgtac aaatccagat ttccatctg tataatttc 11820  
 ctctacctg aaggatgta ttttctc tgtgcaggc ctgttggtga ttaactctt 11880  
 cagcatttt tttttttt tattgacag gatcactct gtcactcagg ctggaatga 11940  
 gtgtacaat tatagttcac tgcagcctg aactcctaga cccaagcaat cctcagcctc 12000  
 ctgagtggt gggactatag gcatgcatca ccacgcctg ataattttt attttactt 12060  
 tttagaga tgggtctc ctacgtgcc caggctgtc tctactct gggctcaagg 12120  
 gatctcca cgagggatcc tccacttg gcttccaaa tgtgggatt acaggtgtga 12180

gccactactc ctggcttct ttaactttt gacttttga tgcctgaaag tctattttgc 12240  
 ctctatttt caaagatagt ttgctgggt atagaattct agacttttt ttcttttca 12300  
 gtactttta aaaaaaagt attattatt tttttatt tgagacagtg tctcgctgtg 12360  
 tcaccaggc tggagtgcag tggcagcgc ttggctcact gcaatctctg cctcctgggt 12420  
 tcaagtgtt ctctgtctc agcctccag gtagatggaa ttacaggcgc atgctaccac 12480  
 gcccggttaa tttgtact ttagcagag atgggggttt gccatgttg ccaggctgt 12540  
 ttgaaacct ctgacctcag ggtgatccgc cctcctggc ctccaaagt gctaggatta 12600  
 caggcgtag ccacttgcc tggcctcagg acttttaaag atgctattct gctgtcttct 12660  
 ggcccactc gttcaagaa gtgtgctggc attcttgggt cctctgtatg taacctttt 12720  
 atccaccgt tttaaatat tcttatcac tggtttagg cagttcata atgtggctg 12780  
 atgtagtgt tttgtttg ttgttttt ttttctgtg ttatgcttg tgatcattg 12840  
 aactcttag acatgtgggt tgcagttc atcaaattg gaaaaaaac acctatctt 12900  
 tctgaaac ttgtactgt cccctatgc gattccacag actttcttg cagaatcccc 12960  
 tgtgtatt tcaggcacct cgaagggtt ctacagctc actgggatt ttcagcctc 13020  
 tctttgtgt gttcatttt ggatagttc tgcgtctaca tttagtctt ttttctgt 13080  
 aatgtctcag ctgtgtaat tgcaccaagt gctgttccg tctcagacac tgtggtttc 13140  
 attctagaa atccagttg aatctttt ttcttttt ttttttta agatggagt 13200  
 tgcctgtt gtccaggctg gagtgcattg gcgcggctt ggctcactg aacctctgc 13260  
 tcccggttc aagtgggtt cctgcctcag cctccagggt agctgggatt ataggcatc 13320  
 accaccatgc ccagctaatt ttgtattt tagtaaagac agagcttcac tgttaggct 13380  
 gggccagtct cgaattcctg acttcagggt attcagctgc ctgcgcctcc caaagtgtg 13440  
 ggattacagg cgtgagctac cgtgccagc ctcaattga atcttttaa atgtatttc 13500  
 tacatctta cttaacagtt gcaatcttc ctctgtttt ttttaagta tatggaatt 13560  
 ggttacagt attctttta tgcctatta actaaattac atttggttac agttactct 13620  
 ttaatgtct gtaattctg tcatcttaa ctatgtact ttgggtctt tttttatt 13680  
 tttatttta ttttgaga cagggtctc ctgtgtacc caggctagag tacagtggca 13740  
 tgatcatggt tcatatagc ctcaacctcc caggctcaag ttatcatct acctagccc 13800  
 ccaagtgc tgggactata ggcatagcc accatgtca actaatttt ttttttgt 13860  
 ggagacagt tctactaat ttgctcaggc tggcctcaa cttctgggt caagcagtc 13920  
 tctgtcttg gcttcccaa gtgtgggat tacaggcatg ggccactgca cccagccata 13980  
 tttctgtct tcttgcatg cctagtaatt ttgtttgga tggatgctag actttgtgag 14040  
 ttgaggta ttgcgtctg gatattttg tattctataa atcttctga gctttgtct 14100  
 gggatacagt taagttact aggacagtc gatccattt ggtctactt ttaagcttg 14160  
 tgagctggac cagagcagt ttactctag ggcaagccta atttgccca ccacttgatt 14220  
 ttgaaataa agtttatg gaaaatcact cccattcatt tacatgttca atgtcaggt 14280  
 ttatgcaata ataacagact tgagtagtg cgccagagac ttatggccc acaaagcaa 14340  
 aagttttag tatttagccc ttacagaaa acatttgcca acccaggcc ttgggcta 14400  
 ctccccact gctgaggtaa aacctctgg agtccctgat gctcatgaa ttatgagggt 14460  
 ttttactct ggtgtgaac ctgggattg tccctgtaa cctctttgt gactctccc 14520  
 cactgtgggt ttgggtcatt ttgcccagc ttgctcga ccagttctt gctgcacac 14580

cgagggcacc tctgcagatc tctgcagctc tgtctccggc tgcctctct gctcctgtga 14640  
ccttgccggc agctccagct gcaggccctc tcagacttc tccagcalt gccccctgaa 14700  
ctcagggaaa cccctgggct ctgctagggg gtcccccca tacacatggg gaccatcggg 14760  
ggaaaccgta ggcagccggc ggggcagtga tagactacc tccctgttt cctgtggcc 14820  
gtcatggctg ctgtcccgcg tcttagtggc catcatgtcc actgtccgc atcttgagg 14880  
ccattgttc tgtgtttg gaaaggggtt aaatctggt tctctcct ttgtgcttg 14940  
agggtgatgt gccctgggcc acctgattc agagagtct tgcgcgggtg cacgatgtcc 15000  
agtcacctgg agcttggcag cagggtggat gtgcacctg ccgcagcctg cagctctgcc 15060  
gtccaccctg ctatgtctg cacatagcat ttatgtctgt gttgtttc ccagagtga 15120  
gagtggagac tgtggcccg tggggcgctc tctgcctct gggatgtctg cccagtaac 15180  
ctggaggcag ccgcgctacc ccacactgc cgggagcagg gtctcggacg cacctcct 15240  
tctctcctag ctgcgaacgc cctgtgggtc tcttcgcag ccctggttc gtactcttc 15300  
gagggtactg gataccagcc ttcatccta acaggggaga cagctgaggg gtcccccca 15360  
gtccgatcc cgcctctc agtgaccaca gccaacggga cgatctcct caccgagatg 15420  
gtgcagggtg gcggagccgg gaggcaggat gccgtggctg aggtgcagt ggccttggc 15480  
ctggctcta cctgatga tctgtgggt gccaggggt ctgaggtcag ttaggacagc 15540  
tgagtctca ggaacggaca tctcagttat taaagaatcc caggltggat gaaactcag 15600  
cgagctcagg gatgtcacgt ttgtttcag gggcgcttct cctgtttg actccagctg 15660  
aggatgaatt taccgttc ctccagcac ctggcgctc ttacagaaag gaggcggatc 15720  
ctgcagctga caagcactg ctctgttac ctgtgggcg ggggtggctc ttgtgctt 15780  
catgggtcac tctgtggctc tacccttag gaaggtcact caccatccct ctctcttc 15840  
tcaggacatg ggagccgggc tggcgtgtt gccctgatg ggcctcctg agagcattgc 15900  
gggtggcaaa gcctcggta agacgcctgt caccacacc ccaggtctcc cagtgcgcg 15960  
gctgggctag gcctgcctg ttctagctt gcctttatcc gtactagt ttgaaaatt 16020  
gaattcatat ccaagtaata catgctcatg atagatacat atgtattgt tatatatg 16080  
aaaactggat ctataatgag gcattgccct ccaccccatg gtgtgtgtt gagtgtta 16140  
acagcctctg ctgtttgtg aaataaaagg ttgtctcg tggccttc cgatgtccac 16200  
gggtgaaacg ctgtgtctg atttaagggt aacgtcactg aaaggggagt ttgcacatg 16260  
agctgggtg agatctgat gaacaatcat attctatgt gtctccacca ttagatata 16320  
gtgggtgcaa ataacctat cagtagtag caaatgcaa ataaattagg aagtgatgag 16380  
tttaagtat tatcttggg ccaggcatg tggccaggc ctgtaatccc aacatttg 16440  
gaggctgaga cgggaggatc gcttgagctc aggagtta aaccaccta ggcaacgta 16500  
cgagacctcg ttctattaa aaataaaat tagctggga tggagcacac ctgtgttacc 16560  
agcttctcag aaggctgagg caggagatc actgagccc gggaggtcga ggtggcattg 16620  
agctgtgata gtgccactg actccagct gggcagcaga acaagatcct gtattatctg 16680  
tttaattgaa agtttaatt taattgtaa taatggctat gtttagtaat aggtcacia 16740  
aactctaaa cattagcaa catgcttat tcagctggct cagccatcg gccagccct 16800  
cctcgcctc agggaggcag ggaaccctc gtctctca gtgtgttt tagtcagca 16860  
tctctagatt cctccttc agctccgtc ttactcgt ccactcctg cagcttga 16920  
ttgtctgc ggaatgacac ttgtctgt ctgtcaccg tcattagtc ctcttgcgt 16980

ttgggtttat gttgattcta aaggttacag cccagttacc agtgtatgtt actgtgactg 17040  
tgtaaatcgt gtttactgcc tagcccaggc tgtgccaggg taaagttcct tctccaaagt 17100  
cccagtgctc taaccttac cccctcaaa ggagttttt ttgtttttt tttttttt 17160  
ttttggaga cagagtctca ctctgtggcc caggctggag tgcagtggca cgaatctctc 17220  
tcactgcaac ctccgcttcc caggtttaag caattctcat gcctcagtct cccaggtagc 17280  
tgggaataca ggcacgcgcc accacgccc actaatattc tatactttta gtagagacag 17340  
ggctcacta tgttggccag gctggctca agtgatctgc ccacctgggc ctccagtaa 17400  
tttttttt gagatagtg ttactcttg ttgccatgc tgggtgcaat 17460  
gggtgatct tggctaccg caacctccg cccccagggt ccagcaatta tctgcctca 17520  
gcctccaag tagctggat tacgggtgtg caccaacatg cctggctaatt ttgtattt 17580  
tagtagagat ggggttctc catgtgatc agactggtct caaactcctg acctcagga 17640  
atccgccac ctacgctcc caaagtgtg ggattacagg tgtagccac catgctcagc 17700  
cctccagtaa ttctttttt tttttttc gagacggagt ctgctctgt cgtccagggt 17760  
ggagtgcagt ggcgtgatct cggctcactg caagctccg ctctgggtt cagccattc 17820  
tctgcctca acctcccgag tagctggac tacaggcgtc cgtccaccag cccagctaatt 17880  
ttttggatt tttagtaaaa acgggggtt caccgggggt ctgcatctcc tgacctggg 17940  
atctgccctg ccttaacctc tcaaagggt gggattacag gctggagccc ctgggccccg 18000  
gccgtaatt tttaatgaa aaacaaggc tcaatttggg ggccaagggt gacctcaaac 18060  
tctgagtc aaggatcct cctgcctcg gttnnnnnn nnnnnnnnn nnnnnnnnn 18120  
nnnnnnnnnn nnnnnnnnn nnnntggtct gggctaatt ttgttttt gtagagagag 18180  
tatttgcat gttgccagg ctggtctga actcctggg tcaagcatc ctctgcctt 18240  
ggccttaaa agtctcgca ttatagctg gagccacct acctggcctg attacatgtt 18300  
actgtacct ttccactct gtgatact caagcacatg ttacacgca catatatgtg 18360  
aatatttag gtgtgattc ttgatttac aaaactatt gccgtgtga gaaatcata 18420  
atctgttta catttttga atacttttg agcagatgtg aatattactg accaggtctt 18480  
taccgttgt tccctgtag atctcagaat aattaccgca tcatgcca ccaggagctg 18540  
ctggccatcg glaagacccc agccgggga aggaagacac cagctgtgg cctccagggt 18600  
cccaggcctg cccctctgt tctctgcat ttaggaaga ccatgatgtt ggtgatgaac 18660  
tgggaggga aagggtggcc cagatgggt ctctggaat atttagttt gtttgggtt 18720  
ttgagatgg agtctcactc tgttggccag gcaggagtgc agtggcacga tctcggtca 18780  
ctacaacctc tgctctctg gttaagtga ttctctgccc tcagcctccc aagtagctgg 18840  
gattaaaggc atatgccacc atgcatggct aatttttga tttgagtag agatgggggt 18900  
tcacatgtt ggccaggctg gtctgaact cctgacctca agtgalccg ccgctcagc 18960  
ctcccaaagt gctgagatta taggcgtgag ccactgcgc tggcctatgc ctgttatct 19020  
taaacctga gactcagat gtggcctgg accagcagcc tgggcactgg gcatctcctg 19080  
ggagcttact ggaaaggcag gatctcaggc cccacccag atctccgaat caggatctg 19140  
attctcagg gcacgcccag gggattcatg gggcagctt agaagtcaaa gtttgaagc 19200  
cgggcatgtt ggctacgcc tgaatcca gcacttggg aggcggagg aggcagatca 19260  
cctgaggtca ggagttgaa accagctgg ccaaacccg tctctactaa aaatacaaaa 19320  
attagctggg tgtgtgtgt tgtcctgt atccagcta ctgggaggt taggcagga 19380

gaatcacttg aacccgggtg ggcggagggt atagttagcc aagattgcac cactgcactc 19440  
cagccctggca acagagcgag actcctctc aaaaaaaaaa aaaaaaagt caaagtttg 19500  
gaaatgcta gagaccccat gttttcaaa gacttgttt gtatgacttt atgagatgat 19560  
caagcagggt tggccacaag gggagacagc caaaggctca ggaagataac cagtttttt 19620  
ttactcaaag gtcccttagt taccagggag gccacagcat gcctacttag gcccalgtca 19680  
gaggacacca ggggtgtccc cggaaggata aatcccaatc tgtgattta ggccctcgcc 19740  
cttttcggg aattttctt gcttaagct aattcaata aaggccccc gcttaagact 19800  
ttcatatgt tattaatttt aggcgggttc taactccta atgaagaaat attcctcct 19860  
ccctggallt ccgaatcaca ccattcttc gagaaacgcc tccacnntnn nnnnnnnnn 19920  
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnngttt tttttggg caccggcca 19980  
tcactgggg aggcagcatt aaggcccatc acttgccct tcaatgtgac tgcactttt 20040  
ttacacgaaa aagtttttg gtgaaaccc ggtggtcggc ccattacct cctgagtgc 20100  
tagggttgt gttccgggcc atttatctca gtttcaccc cctctggat ggaagatgc 20160  
tcccctctgt tctgctggg gaaggcgggt cctggtcca caggatagat tgggtcaaa 20220  
gtgttcaga gctaggatct ctccaggct ctggtgggc actcctgtt cccagcact 20280  
tgaggaggta agccaatgg gtcacctaag gtcaggcatt tgagaccaga ttggccagca 20340  
tggcacactc cctgtgtgt actaaaaata gaaaaattac actgggtgtg gtggcacatg 20400  
cccgtagtcc cagcacttg gcagggtcaa ccgggtggat cactgaggtc aggagttga 20460  
gaccagcctg gccaatatgg caaaactctg tctgtactaa aaatacagaa attagccagg 20520  
cgtgtagtg tgtcctgta atcccagcta ctcaggaggc tgaggcagga gaatcgctg 20580  
aactgggagg cgagagctgc agtgagctga gatcgcgcca ctgcattcca gctgggcaa 20640  
caaagccaga ctctgtca aaaaaaaaaa aaaaaaaaaa aaaggctagg cttacgtct 20700  
gcaagaatgt ggctgtttt ctctctccc tgaaggagt cgtaggcca ggccccagg 20760  
atgattccc cgagccctgc ctctcgtc cctctctgg tttgtccc cctcccatc 20820  
tccttccct ccactctgt tgcctccct ccacgatcag cctgtctgc ctctcccca 20880  
ggctcacca acatgttgg ctccctctc tctctctacc cgttcacagg cagctttga 20940  
cgtgagtga catgtccgc tctctgtt gccacgttg gacgcctaa cgtgttacg 21000  
ctgacaagga gtcgctgc cctgacccc ggcggcgtc ctccactgt aacgtccgt 21060  
ggagaggcag ggctgggggt caccacgt cctccagggt gtctctgt tcttattct 21120  
catagatcgt cctgcagtt catactagaa agttccact ggcatgttg tacaccctg 21180  
ttatccatt tactctggag gcggaagcag gagaatcgt tgagccagc aggatgaggc 21240  
tgcagttagt tgtaccgca ccgtgcact ccagcctgg caacagagcg agaccctgc 21300  
tctaaataa atactagaaa gtcccagca cgcaaagcc ctctagctc ctggtgccag 21360  
agtcaattcc tgaaaggacg tgagatagg aaggccctc gctctgtc gaagcagcg 21420  
ggcatgaagc ttgcccaga tgcctacgg cccctctca gtcaggaca caggatggag 21480  
gtgacctgt gctaaagga gagaaggagg cgtcgcttg cactgccag tccccagct 21540  
ggtgacctt gccctgtg gtatggggc cccacctga tggggcagg agacagatc 21600  
ggcaggaacc tgaaaggaca cgtgctcct gagctctc ctatagtcag ggtggccaa 21660  
gcgcggtgt ctgtactgc accctaagtc tcttgctc ggtccctg cagtccccg 21720  
ctgctcca agcgtgtg ggagctgacg tccctcga agatcagca caggagtgtg 21780

gactgaggtc tccctttcc cggccctgg tgactgacgg tctctgtgt gcctccagg 21840  
 acagccgtga acgtcagtc ggggggtgc accccggcgg ggggcctggt gacgggtaag 21900  
 gcccccatc tccccttgt gccgcagcc ctgagagtgg gagaaagga ggagggggcc 21960  
 cacagagacg tcccttggc tcatgggccc tgcgccccg gactgcacag ggacttggg 22020  
 ggccacacag gagtagggg accacaggag actgagcagg ggcgggggc ctggcagtc 22080  
 gtgccttac cccaccct glcccagtg ggctctgtg aacaagagg tgctacgtg 22140  
 cgtgtgggg ggacctgca ctccgaggt cactgtgt cccgtcccc gcaggagtgc 22200  
 ttgtctgtgt gtctggac tacctgacct cactgtcta ctacatccc aagtctgcc 22260  
 tggctgccgt catcatcatg gccgtggccc cgtgttga caccaagatc ttcaggacgc 22320  
 tctggcgtgt taagagtacg tcctgtct acaggggaga gcgtgtgat gcggtgtctg 22380  
 aacggaggag gtgtcattt tgctaccca tttctgca gccccctg tggggctggg 22440  
 actgggaagt tagggcagtc ccggaacaga gaagtggatg gccaggagat gggccagag 22500  
 atgtlccga ggctcagtg gaagagctgg agctcctgt cctgacacct ggggtctga 22560  
 ggcgagcact gaccggggg agggccccct cctgatccc ctgccccat cctaccctc 22620  
 ctggcaccg gccccagcc accactctgc ccggcccagc tggggggagg gacaggagac 22680  
 gtccctggg accagcagg ccagcggaa agcctgcac cctggctcag aatggcagtt 22740  
 cctttttt ttattatt tttttatt tttttatt gatcattct ggggtttct 22800  
 tgcagagggg gatttggcag ggcatagga caatatggg ttgggggtaa ggtcacagat 22860  
 aacaggatcc caaggcagag gaattttct tagtcagaa caaatgaaa agtclccat 22920  
 gtctactct ttctacacg acacagcaac catccgatt ctcaatctt tccccactt 22980  
 tccccctt ctattccaca aaaccgcat tgcatcatg gcgcgtctc agtgagctgt 23040  
 tgggtacacc tccagacgg ggtggggcc gggcagagg gctcctcact tccagtagg 23100  
 ggcggccggg cagaggcgc cctcacctcc cggacgggag cagtctctt aactaacac 23160  
 atttgttt gttgtggaa gaagcatgga gtgggtcac ctacttact tggcatttc 23220  
 tggtagccc cagcaagtg ccaggtagaa ggaaggataa tttctgca tgccgctgc 23280  
 atgccagggc ctttcgctg ccagtgcgt gggccttag aacagccctg aaacgtgtc 23340  
 atgtccggc gtggaatgg atggccgctg ctgctagaaa ccaaggctca gccagggtg 23400  
 ggcaggcctg acctatcgt gcggtggaat cccccacagg gctaagcccg tgcactttgt 23460  
 cccaggga cctctcct ggccaggct caagggtca cgtgtccct gcccactcc 23520  
 tcaggccagc tctgtccct gacaagcccc tgctgtgcc ctccctagg ttggaggcca 23580  
 gggaccggc gccaggtct accgtgctc agtcagatg gggagggcat tctttctt 23640  
 cgactgaag catggcctg tcagcagctg ctgtcccaa gtctcaggg gctgcttgg 23700  
 gtccatgag acctttac atagtggg ggcagaagg tgtccgctg tgcagaggg 23760  
 ccatgtggg gcccgggc gctccggg attcctcagc tgtccctc tctagggt 23820  
 ggacctgtg cccctgtcg tgacctct gctgtctc tggagggtc agtacggcat 23880  
 cctggccggg gccctgtgt ctctgtcat gctctgcac tctcagcca ggcctgagac 23940  
 caaggtaccc ctccgtggc tctagtgg gagtgtctg ggggcaggat tctgggcat 24000  
 ggtctatgt ttgagggc cgggtgatt gtgtcgtg gtgtctga aggggaccg 24060  
 tcgtggcag gtgggcagc acctgtct aaacctggt gttctccac tgttggggg 24120  
 ccgtgggggt cccccctg caccctc cgggtccct gcagcacgt aggtgggtg 24180



ggggctccc gcttgggaca ggccaagcct ggtggaggcc acccggtcag acccgctcc 24240  
aggactcact cctccccaca ggtgtcagag gggccgggtc tggctctgca gccggccagc 24300  
ggcctgtcct tccctgccat ggaggctctg cgggaggaga tccaaagccg ggccctggaa 24360  
ggtgcatggg cgggggtcaa ggtggtctga ggtcactccc ctgtctctg cccccactc 24420  
cctgtgttc aggaccccaa gacctgtcc ccgacgtct ccagtcaca aggatgcagg 24480  
catctctgag tgggtggac cgtctctgt gggcctcagc cagtggctgc tgcagcaagg 24540  
gtgtggctc cccacatac actcctccc tggccctaaa gtccggtcc tgttctggg 24600  
gggttgatt taggggagct aagggcctgt ggtcctagg agggaaacag ctgtgtctgt 24660  
caccaaaca tttctctgg tctgccacc cgaatcccc aactgggca ctcagccgc 24720  
acgagatgga gcactctggc ctctgtcc cctgccctg gccagagcct ctttggcct 24780  
ctgcagagca gcttgggt gctctgggt gctgacctg gctgggct gctcccccag 24840  
tgtccccgc acgtgcctg tctctggagt gcacccatgt ctgcagcatc gactacactg 24900  
tgggtctgg actcggcag ctctccagg actccagaa gcagggcgc gccctggcct 24960  
ttgtggcct gcagggtgggt gtgcactgg atgcctagg ggtagcagc tgcggaagg 25020  
ccttctgtg cctgcctccc atggcgaatg tgacatctt gggctgtat gctggacgc 25080  
cctcggcgc gtgtggctc tctctgat taaacagtt ctgtccca tctggcctc 25140  
ctgtccctc cctgtgaag ggggagcgg gtccccagc cctccaggg gtcagttat 25200  
ggcttctgt cactgccaca ctgtccttg tctggggac acacagtga cgagggcag 25260  
tccctgtct cagggaatt gtaattagg aaacaaaag agacaggtga ggtaggcgt 25320  
ggtggcacac tctggaatc ccagcactt gggaggcaa ggcgggcaga tcacctagg 25380  
tcaggagtc aagaccagc tggccaacat ggtgaaacc cgtctact aaaaataca 25440  
aaaaaactag tcaggtgtg tggcgagcg ctgtaattc agctacatg gaggtggga 25500  
gaatcgctt aacctgggag gtggaggtg cagtaagcc agatccact actgtactc 25560  
agcctggcg acagagtgg actccatc aaaaacaac aaacgaaaa aaacaggtc 25620  
tcatagaatt tcatgaaaa cgtatttca gggctccag aggtgaaga cgggttcta 25680  
tggaggcgt cctgtcaga gccgcagga aagtgaagg gctgggtcc aggcctgcg 25740  
tctaggcct cacctaggag cctctgagc actgcagggt caacatcca ggggtgtggc 25800  
cagtgttc agagaggcag ggtctctgt tctgtggt aatgtgcg tcttgcca 25860  
gtcctagat gcagaacgt ggggactagg gctgtccc aggcgccag aggagacatt 25920  
catgaactag ccatggaaca ggaggccaag tgaccgtat acccagggtg tggaccacag 25980  
ccgaccttg tcagagttc ctctctcc aacctggcc acgaggctag tttatctc 26040  
atccccatg ccagaagagg tcactaggc ccagagaagc caggaatct gccacggc 26100  
acacaggatg gtgggccag ctctcagatt ggcctgtgc ctcaagaact tgcctcata 26160  
gtgtacgc ctgtcata ttgcagagc acatgtct atgtcttg gagatgatt 26220  
actaaaaa atactctct gaccagaca tggtgctca cacctgta cccagcatt 26280  
tgagaggccg aggcgggagg tcaagggtt gagaccagc tggccaacat agtgaactc 26340  
cgtctact aaaaataca aaattagac ggcattgtg cgcacgctg taatccagc 26400  
tgctcaggg gctgaggcag gagaatcgt tgaaccggg aggcagaggt tgcagtgc 26460  
cgagatcgc ccattgcact ccagcctgg taacaagag gaaactccat ctcaaaaaca 26520  
aacaacaaa caaacaaaa gtccttgg ccaggcgctg tggctcacac ctgtaatcc 26580

agcactttgg gaggccgagg taggaggatc attgaggcc aggagttga gaccagcctg 26640  
 ggcaacatga cgaaccccca tcaaaaaaat aggaaaaaat agctggacgt ggtggtgtgc 26700  
 acctgtagct actcaggagg ctgagggtgg aggatcactt gactctggga ggtttaggct 26760  
 gcagttagcc atgatcgtc cactgcactc cagcctgagc aatgagcaag accctgtctc 26820  
 aaaaaacaaa aaatttaaaa aaaaaaaatc ttctctgac agcattcccc tggggctgcg 26880  
 ttctctca ccattcactg gtatggagggt gaagccatac ctctccggga gactctgaga 26940  
 .tggcatgtct accaggctgc cgaccctgt gctacagagg aacatccctg ccctggctaa 27000  
 agtctgtctg tctctcagggt ccccgctctc cgtgtcctgc tgtccgtga cctgaagggg 27060  
 ttccagtact tctctacctt ggaagaagca ggtgggcaca gtcagacatc ctgtggcttt 27120  
 ggtgattttg taaaaatcat aaatgcttat tgaaaaaat atgggaaacc agctgggcac 27180  
 agtggctcat gcctgtaac ccagcacttt gggaggccga ggcagggtga tcacctgagg 27240  
 tcaggagttc gagaccagcc tggccaacat ggtaaacct catcagtact aaaaatacaa 27300  
 aaattagctg ggtgtggtg catgcacctc tagtcccagc tattcgggag gctgaagcag 27360  
 cagaatcgct tgaaccagg aggagagggt tgcactgagc cgagattgtg cctctgact 27420  
 tccagagtag ggtgagagag ctagccatgt tgtccagaaa annnnnnnnn nnnnnnnnn 27480  
 nnnnnnnnn nnnnnnnnn nnnnnnnnn ncttgggtga gcaggataa tgccttcaa 27540  
 cccaggtagc agcagttgta ctgagtcga gaattgtcc actgcactcc agcctgggtg 27600  
 agagcagcga gtactctgc tcaaaaaaaa aaaaaacaaa aaaaacagga aaccacctcc 27660  
 accccagtc actttgtga tcaactcata cccctccct aaacacgcat atgtacctgc 27720  
 ctgtcaggat gtggatatat gtttgcggt ttacgtaaat gggaccattt calacctgt 27780  
 gctctggaac ccacatttt catcagaag gttggaagga tgccttcca gccgaaagtc 27840  
 cacatcccg gggatcagga cagagcagg cgggtcagg agatccagaa gcccgggac 27900  
 agaaggtagc ggaggacag gagcagggt ggcgctgacc cttgagacag caatcgagg 27960  
 aggtctgagc cgcagcagg gtcaacaaga ggaaggcca gagatgcaga gcacccacc 28020  
 caggccacac agcagtgcc agagggctcc agggccaggt gctaggctc ttctcttc 28080  
 actgaggtca cagctgaagc tgggtcagct ccgtgagagt gagggtggc ggaattgta 28140  
 ctgacttct tggctcaatg tgacgtcagg gagatcacc catgtgttg tagaatcagc 28200  
 tcagatgcag tgcactggag gtgataagc agaattgtgc ctggcgtg gatgggagg 28260  
 taccctggag cagtaagaaa gggccgttag tcacctgaaa aatacgctta cagagactca 28320  
 ggtgagacc tcattggagt agtgacatc gccgtgggtg ccacagctc ctctctcac 28380  
 acctccagg actctggaag gccctctta atccctct gtgaactgac ccactctac 28440  
 ttctgagct ttagtgctg aaacattat tgtatttct gcagagaagc acctgaggca 28500  
 ggagccaggg acccagccct acaacatcag agaagactcc attctggacc aaaagggtgc 28560  
 cctgtcaag gcataatgg gccacccgtg ggcacccaca gtttcagggt tgtccgga 28620  
 ggttctgtc actgtgatt gatgtggat gccgctgat agacatgtc gcctggctga 28680  
 gaaacccctg agcaggaac ccagggaaga gaaggagcc aggcctggag gtccacggca 28740  
 gtggggagtg gggctcactg gcttctgtg ggaatgactg aaaatgacct cgtgtctgt 28800  
 ccctggcatg accctcttg gaagagtggt ttggagagag cctctagaa tgacagactg 28860  
 tgcaggaag caggggcagg ggttccagc ccgggctgtg cgaggcatcc tggggctgc 28920  
 agcaccttc cggctacca gtgccactg cgggggagg acggggcagg caggagtctg 28980

ggaggcgggt ccgcctctct tctctcggc atctgtgctc tccgagagaa aaccaagggt 29040  
tgtcaaatga cgtcaagtct ctatttaaaa ataattttgt ttttctaaa tggaaaaagt 29100  
gatagctttg gtgattttgt aaaagtcata aatgcttatt gtaaaaaata caggaaacca 29160  
ccccctcccc tgtccacttg ggtgatcatt ccagaccctt ccccaaacat gcatatgtac 29220  
ctgtccgtca gtgtgtggat gtatgtttac agttctacat aaatgggac atttataca 29280  
tgggtgcttg gaaccacat tttcatgca gtcatgtgca gtgaattatt tattgtgata 29340  
ataaatagca ttagaataca agattttta tctctcgtg gtattttgt ctatatatgc 29400  
atcataattg acttaccgag cctctgttc aacgtgtgcg tgggttagag acgggatcgt 29460  
gcctccttta gatgtgtcag tgaagccct tgcctctat gtgactgcgt ttgaaacag 29520  
ggcttttagg gggtaactag gtttaatga ggtcataaag gtggccctg aaccgatagc 29580  
ttagtgtcc tctgagaag agacgccaga gagctcgtt tctgaccct cacacccag 29640  
ggaactccat gagaggacac ggcaaaacca ggcatgcgc ccaccaggaa gaaaggcctc 29700  
atgaggaccc cagcctcaa aactgggaga agatgaatct ctgtgctagg cccgcagcc 29760  
tggggtgatc tgtacggcg gcctgagcag gtgaggactg cctgcatgtt tgtatttta 29820  
tgaatgcttt gattgagctt gggggtaa at cctggaggc ctgtggcagc ctacagggtg 29880  
tgtctccct gcacttctt caagaagaat gtggcctgcc ctgctgagcc tctgtctgcc 29940  
cgttctgcc gggcagtc ccgaatgtc agcgcagcaa ggggagggcc tctgaacca 30000  
gggctgtcgg ctgctgggct cccactgga caggggagcg gacattggag tgccttcat 30060  
ccgctgact ctccaggtc cctgcctcc catttctt ccttcttcc tcttcttcc 30120  
cttcttctt tcttcttctt gctccttcc ttttcttct tcttcttctt ttttcttct 30180  
ttcttcttct ttttcttctt tcttcttcc tcttcttctt ccttcttctt 30240  
cttctcactg tgttctccag gctgttctt aactcctggg ctcaagcaat cctctgcct 30300  
cagcctctgg tgtggctgg accacagaca catgccaccg cgcaggcta attgttaat 30360  
tttttata gagacagggt ctacttgtt tcccaggct cgtctcaaac tctgggctc 30420  
aagtgtact ccagctcag cctgaagtgc tgaattaag gtgtgagcca ccatgcctg 30480  
ccctcttca ttcacaagt aacgttcac cctgcctc caatccatgt cgttctgac 30540  
ctcaggcagc tttcttctt acataaagt aacctgtccc aaagctgtgc tactgggcc 30600  
ccccgccag ggttgaaga ggcagcagtt cacattggc ttgactcac ataccaagg 30660  
catccatgct tgaacctaga catggttcat tcacaggat ggggaagg aaacagactg 30720  
tggcgggcac taggcactat cagttcata aatctgata ccactgtga ctacagcgtt 30780  
atgcttctg gaatctact aaacatgctt gctcagcct tccaggaagc aggcacaagg 30840  
acgtttgtga caactggc agtaattta agaagtagga agaaacctga gtttctga 30900  
attcggtaac atctgactg taggacacac gaataatgcc gtggaatait gtgcagcagt 30960  
taaaagaaat gagggaagg cgggtgtgg cagctcacgc ctgtatctc agcacttgg 31020  
gaggccgagg ctgggtgat acttaaggc aggagttga gaccagtctg gccaacatgg 31080  
cgaaacctg tcttactaa aaalacaaa attagctgac ctctgtgaca gagtgagact 31140  
ctgcctcaa aaaaaaaaaa aaaaaaaatgagg tgaagcttta agttgaaca 31200  
ctgatttgg ggtgcaata agcaagtgc agaataat ccatgttaag atgctattg 31260  
agtgaacaca cggaccaaac aattctatgt tgggtacaaa ttaagaaag agttctggc 31320  
tgggcacggg ggtcacaaac tgtaatcca gcactctgg agggtagtc ggaatgatca 31380

ggaggtcagg actctagac tagcctggcc aatatggtga aaccccatct ctactaaaaa 31440  
 tacaaaaatt agccaggcgt gttggcgcgt gctggagtc ccagctactc aggagactaa 31500  
 ggtaggagaa tcactgaac ccgggaggca gaggttcag tgagctgaga tcatgccact 31560  
 gcactctggc ctgggcaaca gagcgagact caaaaaataa ataaataaat aaataaaata 31620  
 aataagagtt ttctgcactt tgggaggcct gtagtcccag ctactctgga ggctgaggca 31680  
 agaggatcac ttgagcctgg gggggtcgag gctgcagtga gtctgattg tgcactgaa 31740  
 atccagcctg ggcaacagag tgagac 31766

<210> 6

<211> 548

<212> PRT

<213> Human

<400> 6

Cys Arg Pro Ser Thr Val Thr Asn Lys Phe Pro Ile Leu Lys Trp Leu  
 1 5 10 15  
 Pro Arg Tyr Arg Leu Glu Tyr Ile Met Gln Asp Phe Ile Ala Gly Phe  
 20 25 30  
 Thr Val Gly Leu Thr Thr Ile Pro Gln Ala Ile Ala Tyr Gly Val Val  
 35 40 45  
 Ala Gly Leu Glu Pro Gln Tyr Gly Leu Tyr Ser Ala Phe Met Gly Cys  
 50 55 60  
 Phe Thr Tyr Ile Val Phe Gly Ser Cys Lys Asp Val Thr Ile Ala Thr  
 65 70 75 80  
 Thr Ala Ile Met Ala Leu Met Val Asn Gln Tyr Ala Thr Ile Ser Pro  
 85 90 95  
 Asp Tyr Ala Val Leu Val Cys Phe Leu Ala Gly Cys Ile Val Leu Leu  
 100 105 110  
 Leu Gly Leu Leu Asn Met Gly Val Leu Val Arg Phe Ile Ser Ile Pro  
 115 120 125  
 Val Ile Thr Gly Phe Thr Met Ala Ala Ala Thr Thr Ile Gly Ser Ala  
 130 135 140  
 Gln Ile Asn Asn Ile Val Gly Leu Thr Ser Pro Ser Asn Asp Leu Leu  
 145 150 155 160  
 Pro Ala Trp Lys Asn Phe Phe Thr His Leu Thr Ser Ile Arg Leu Trp  
 165 170 175  
 Asp Ala Leu Leu Gly Val Ser Ser Leu Val Phe Leu Leu Leu Met Thr  
 180 185 190  
 Arg Val Lys Asp Ile Lys Trp Gly Asn Arg Ile Phe Trp Lys Tyr Leu  
 195 200 205

Gly Leu Ser Arg Asn Ala Leu Ala Val Ile Phe Gly Thr Phe Leu Ala  
 210            215            220  
 Tyr Ile Leu Ser Arg Asp Gly Asn Gln Pro Phe Arg Val Thr Gly Asn  
 225            230            235            240  
 Ile Thr Ala Gly Val Pro Pro Phe Arg Leu Pro Pro Phe Ser Thr Thr  
 245            250            255  
 Val Asp Gly Glu Tyr Val Ser Phe Gly Glu Met Ile Ser Thr Val Gly  
 260            265            270  
 Ala Ser Leu Gly Ser Ile Pro Leu Ile Ser Ile Leu Glu Ile Val Ala  
 275            280            285  
 Ile Ser Lys Ala Phe Ser Lys Gly Lys Ile Val Asp Ala Ser Gln Glu  
 290            295            300  
 Met Val Ala Leu Gly Met Cys Asn Ile Met Gly Ser Phe Val Leu Ser  
 305            310            315            320  
 Met Pro Val Thr Gly Ser Phe Thr Arg Thr Ala Val Asn Asn Ala Ser  
 325            330            335  
 Gly Val Lys Thr Pro Leu Gly Gly Ala Val Thr Gly Ala Leu Val Leu  
 340            345            350  
 Met Ala Leu Ala Phe Leu Thr Gln Thr Phe Tyr Phe Ile Pro Lys Cys  
 355            360            365  
 Thr Leu Ala Ala Ile Ile Ile Ala Ala Met Ile Ser Leu Val Glu Leu  
 370            375            380  
 His Lys Ile Lys Asp Met Trp Lys Ser Lys Lys Lys Asp Leu Phe Pro  
 385            390            395            400  
 Phe Val Val Thr Val Leu Thr Cys Met Phe Trp Ser Leu Glu Tyr Gly  
 405            410            415  
 Ile Leu Cys Gly Ile Gly Ala Asn Met Val Tyr Ile Leu Tyr Ser Ser  
 420            425            430  
 Ala Arg Pro His Val Asp Ile Lys Leu Glu Lys Ile Asn Gly His Glu  
 435            440            445  
 Val Ser Val Val Asp Val Lys Gln Lys Leu Asp Tyr Ala Ser Ala Glu  
 450            455            460  
 Tyr Leu Lys Glu Lys Val Val Arg Phe Leu Asn Asn Gln Asn Gly Glu  
 465            470            475            480  
 Thr Gln Leu Val Val Ile Lys Gly Glu Glu Ile Asn Ser Ile Asp Tyr  
 485            490            495  
 Thr Val Ala Met Asn Ile Val Ser Met Lys Gly Asp Leu Glu Ala Leu  
 500            505            510  
 Asn Cys Ala Met Ile Cys Trp Asn Trp Asn Ile Ala Ser Ala Gly Val  
 515            520            525

Val Cys Arg Leu Asn Asn Asp Leu Arg Pro Ile Phe Lys Phe Asp Leu .  
 530            535            540  
 Ser Leu Glu Glu  
 545

<210> 7

<211> 548

<212> PRT

<213> Drosophila melanogaster

<400> 7

Cys Arg Pro Ser Thr Val Thr Asn Lys Phe Pro Ile Leu Lys Trp Leu  
 1            5            10            15  
 Pro Arg Tyr Arg Leu Glu Tyr Ile Met Gln Asp Phe Ile Ala Gly Phe  
 20            25            30  
 Thr Val Gly Leu Thr Thr Ile Pro Gln Ala Ile Ala Tyr Gly Val Val  
 35            40            45  
 Ala Gly Leu Glu Pro Gln Tyr Gly Leu Tyr Ser Ala Phe Met Gly Cys  
 50            55            60  
 Phe Thr Tyr Ile Val Phe Gly Ser Cys Lys Asp Val Thr Ile Ala Thr  
 65            70            75            80  
 Thr Ala Ile Met Ala Leu Met Val Asn Gln Tyr Ala Thr Ile Ser Pro  
 85            90            95  
 Asp Tyr Ala Val Leu Val Cys Phe Leu Ala Gly Cys Ile Val Leu Leu  
 100            105            110  
 Leu Gly Leu Leu Asn Met Gly Val Leu Val Arg Phe Ile Ser Ile Pro  
 115            120            125  
 Val Ile Thr Gly Phe Thr Met Ala Ala Ala Thr Thr Ile Gly Ser Ala  
 130            135            140  
 Gln Ile Asn Asn Ile Val Gly Leu Thr Ser Pro Ser Asn Asp Leu Leu  
 145            150            155            160  
 Pro Ala Trp Lys Asn Phe Phe Thr His Leu Thr Ser Ile Arg Leu Trp  
 165            170            175  
 Asp Ala Leu Leu Gly Val Ser Ser Leu Val Phe Leu Leu Leu Met Thr  
 180            185            190  
 Arg Val Lys Asp Ile Lys Trp Gly Asn Arg Ile Phe Trp Lys Tyr Leu  
 195            200            205  
 Gly Leu Ser Arg Asn Ala Leu Ala Val Ile Phe Gly Thr Phe Leu Ala  
 210            215            220

Tyr Ile Leu Ser Arg Asp Gly Asn Gln Pro Phe Arg Val Thr Gly Asn  
 225            230            235            240  
 Ile Thr Ala Gly Val Pro Pro Phe Arg Leu Pro Pro Phe Ser Thr Thr  
               245            250            255  
 Val Asp Gly Glu Tyr Val Ser Phe Gly Glu Met Ile Ser Thr Val Gly  
               260            265            270  
 Ala Ser Leu Gly Ser Ile Pro Leu Ile Ser Ile Leu Glu Ile Val Ala  
               275            280            285  
 Ile Ser Lys Ala Phe Ser Lys Gly Lys Ile Val Asp Ala Ser Gln Glu  
               290            295            300  
 Met Val Ala Leu Gly Met Cys Asn Ile Met Gly Ser Phe Val Leu Ser  
 305            310            315            320  
 Met Pro Val Thr Gly Ser Phe Thr Arg Thr Ala Val Asn Asn Ala Ser  
               325            330            335  
 Gly Val Lys Thr Pro Leu Gly Gly Ala Val Thr Gly Ala Leu Val Leu  
               340            345            350  
 Met Ala Leu Ala Phe Leu Thr Gln Thr Phe Tyr Phe Ile Pro Lys Cys  
               355            360            365  
 Thr Leu Ala Ala Ile Ile Ile Ala Ala Met Ile Ser Leu Val Glu Leu  
               370            375            380  
 His Lys Ile Lys Asp Met Trp Lys Ser Lys Lys Lys Asp Leu Phe Pro  
 385            390            395            400  
 Phe Val Val Thr Val Leu Thr Cys Met Phe Trp Ser Leu Glu Tyr Gly  
               405            410            415  
 Ile Leu Cys Gly Ile Gly Ala Asn Met Val Tyr Ile Leu Tyr Ser Ser  
               420            425            430  
 Ala Arg Pro His Val Asp Ile Lys Leu Glu Lys Ile Asn Gly His Glu  
               435            440            445  
 Val Ser Val Val Asp Val Lys Gln Lys Leu Asp Tyr Ala Ser Ala Glu  
               450            455            460  
 Tyr Leu Lys Glu Lys Val Val Arg Phe Leu Asn Asn Gln Asn Gly Glu  
 465            470            475            480  
 Thr Gln Leu Val Val Ile Lys Gly Glu Glu Ile Asn Ser Ile Asp Tyr  
               485            490            495  
 Thr Val Ala Met Asn Ile Val Ser Met Lys Gly Asp Leu Glu Ala Leu  
               500            505            510  
 Asn Cys Ala Met Ile Cys Trp Asn Trp Asn Ile Ala Ser Ala Gly Val  
               515            520            525  
 Val Cys Arg Leu Asn Asn Asp Leu Arg Pro Ile Phe Lys Phe Asp Leu  
               530            535            540

**WO 02/059306**

**PCT/US01/42809**

Ser Leu Glu Glu  
545